

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/349846558>

# Certain new subclasses of $m$ -fold symmetric bi-pseudo-starlike functions using $Q$ -derivative operator

Article · March 2021

DOI: 10.30538/psrp-oma2021.0081

---

CITATIONS

3

READS

310

1 author:



[Timilehin Gideon Shaba](#)

Landmark University

57 PUBLICATIONS 369 CITATIONS

SEE PROFILE

Article

# Certain new subclasses of $m$ -fold symmetric bi-pseudo-starlike functions using $Q$ -derivative operator

Timilehin Gideon Shaba

Department of Mathematics, University of Ilorin, P. M. B. 1515, Ilorin, Nigeria.; shabatimilehin@gmail.com

Communicated by: Absar ul Haq

Received: 17 September 2020; Accepted: 21 February 2021; Published: 28 February 2021.

**Abstract:** In this current study, we introduced and investigated two new subclasses of the bi-univalent functions associated with  $q$ -derivative operator; both  $f$  and  $f^{-1}$  are  $m$ -fold symmetric holomorphic functions in the open unit disk. Among other results, upper bounds for the coefficients  $|\rho_{m+1}|$  and  $|\rho_{2m+1}|$  are found in this study. Also certain special cases are indicated.

**Keywords:**  $m$ -fold symmetric bi-univalent functions, analytic functions, univalent function.

**MSC:** 30C45.

## 1. Introduction

Let  $\mathcal{A}$  be the family of holomorphic functions, normalized by the conditions  $f(0) = f'(0) - 1 = 0$  which is of the form

$$f(z) = z + \rho_2 z^2 + \rho_3 z^3 + \dots \quad (1)$$

in the open unit disk  $\Omega = \{z; z \in \mathbb{C} \text{ and } |z| < 1\}$ . We denote by  $\mathcal{G}$  the subclass of functions in  $\mathcal{A}$  which are univalent in  $\Omega$  (for more details see [1]).

The Keobe-One Quarter Theorem [1] state that the image of  $\Omega$  under all univalent function  $f \in \mathcal{A}$  contains a disk of radius  $\frac{1}{4}$ . Hence all univalent function  $f \in \mathcal{A}$  has an inverse  $f^{-1}$  satisfy  $f^{-1}(f(z))$  and  $f(f^{-1}(v)) = v$  ( $|v| < r_0(f)$ ,  $r_0(f) \geq \frac{1}{4}$ ), where

$$g(v) = f^{-1}(v) = v - \rho_2 v^2 + (2\rho_2^2 - \rho_3)v^3 - (5\rho_2^3 - 5\rho_2\rho_3 + \rho_4)v^4 + \dots \quad (2)$$

A function  $f \in \mathcal{A}$  denoted by  $\Sigma$  is said to be bi-univalent in  $\Omega$  if both  $f^{-1}(z)$  and  $f(z)$  are univalent in  $\Omega$  (see for details [2–11]).

A domain  $\Psi$  is said to be  $m$ -fold symmetric if a rotation of  $\Psi$  about the origin through an angle  $2\pi/m$  carries  $\Psi$  on itself. Therefore, a function  $f(z)$  holomorphic in  $\Omega$  is said to be  $m$ -fold symmetric if

$$f\left(e^{\frac{2\pi i}{m}} z\right) = e^{\frac{2\pi i}{m}} f(z).$$

A function is said to be  $m$ -fold symmetric if it has the following normalized form

$$f(z) = z + \sum_{\phi=1}^{\infty} \rho_{m\phi+1} z^{m\phi+1} \quad (z \in \Omega, \quad m \in \mathcal{N} = \{1, 2, 3, \dots\}). \quad (3)$$

Let  $\mathfrak{S}_m$  the class of  $m$ -fold symmetric univalent functions in  $\Omega$ , that are normalized by (3), in which, the functions in the class  $\mathfrak{S}$  are one-fold symmetric. Similar to the concept of  $m$ -fold symmetric univalent functions, we introduced the concept of  $m$ -fold symmetric bi-univalent functions which is denoted by  $\Sigma_m$ . Each of the function  $f \in \Sigma$  produces  $m$ -fold symmetric bi-univalent function for each integer  $m \in \mathcal{N}$ .

The normalized form of  $f(z)$  is given as in (3) and the series expansion for  $f^{-1}(z)$ , which has been investigated by Srivastava *et al.*, [12], is given below:

$$\begin{aligned}
 g(v) &= f^{-1}(v) \\
 &= v - \rho_{m+1}v^{m+1} + [(m+1)\rho_{m-1}^2 - \rho_{2m+1}]v^{2m+1} \\
 &\quad - \left[ \frac{1}{2}(m+1)(3m+2)\rho_{m+1}^3 - (3m+2)\rho_{m+1}\rho_{2m+1} + \rho_{3m+1} \right].
 \end{aligned}
 \tag{4}$$

Some of the examples of  $m$ -fold symmetric bi-univalent functions are

$$\left\{ \frac{z^m}{1-z^m} \right\}^{\frac{1}{m}},$$

$$[-\log(1-z^m)]^{\frac{1}{m}},$$

and

$$\left\{ \frac{1}{2} \log \left( \frac{1+z^m}{1-z^m} \right)^{\frac{1}{m}} \right\}.$$

For more details on  $m$ -fold symmetric analytic bi-univalent functions (see [5,12–17]).

Jackson [18,19] introduced the  $q$ -derivative operator  $\mathcal{D}_q$  of a function as follows;

$$\mathcal{D}_q f(z) = \frac{f(qz) - f(z)}{(q-1)z}
 \tag{5}$$

and  $\mathcal{D}_q f(0) = f'(0)$ . In case of  $g(z) = z^k$  for  $k$  is a positive integer, the  $q$ -derivative of  $f(z)$  is given by

$$\mathcal{D}_q z^k = \frac{z^k - (zq)^k}{(q-1)z} = [k]_q z^{k-1}.$$

As  $q \rightarrow 1^-$  and  $k \in \mathcal{N}$ , we get

$$[k]_q = \frac{1-q^k}{1-q} = 1 + q + \dots + q^{k-1} \rightarrow k,
 \tag{6}$$

where ( $z \neq 0, q \neq 0$ ). For more details on the concepts of  $q$ -derivative (see [5,20–27]).

**Definition 1.** [28] Let  $f(z) \in \mathcal{A}$ ,  $0 \leq \chi < 1$  and  $\sigma \geq 1$  is real. Then  $f(z) \in L_\sigma(\chi)$  of  $\sigma$ -pseudostarlike function of order  $\chi$  in  $\Omega$  if and only if

$$\Re \left( \frac{z[f'(z)]^\sigma}{f(z)} \right) > \chi.
 \tag{7}$$

Babalola [28] verified that, all pseudostarlike function are Bazilevic of type  $(1 - \frac{1}{\sigma})$ , order  $\chi^{\frac{1}{\sigma}}$  and univalent in  $\Omega$ .

**Lemma 1.** [1] Let the function  $\omega \in \mathcal{P}$  be given by the following series  $\omega(z) = 1 + \omega_1 z + \omega_2 z^2 + \dots$  ( $z \in \Omega$ ). The sharp estimate given by  $|\omega_n| \leq 2$  ( $n \in \mathcal{N}$ ) holds true.

In [29] Girgaonkar *et al.*, introduced a new subclasses of holomorphic and bi-univalent functions as follows:

**Definition 2.** A function  $f(z)$  given by (1) is said to be in the class  $\mathcal{M}_\Sigma(\chi)$  ( $0 < \chi \leq 1, (z, v) \in \Omega$ ) if  $f \in \mathcal{E}$ ,  $|\arg(f'(z))^\sigma| < \frac{\chi\pi}{2}$  and  $|\arg(g'(v))^\sigma| < \frac{\chi\pi}{2}$ , where  $g(v)$  is given by (2).

**Definition 3.** A function  $f(z)$  given by (1) is said to be in the class  $\mathcal{M}_\Sigma(\psi)$  ( $0 \leq \psi < 1, (z, v) \in \Omega$ ) if  $\vartheta \in \Sigma$ ,  $\Re[(f'(z))^\sigma] > \psi$  and  $\Re[(g'(v))^\sigma] > \psi$ , where  $g(v)$  is given by (2).

In this current research, we introduced two new subclasses denoted by  $\mathcal{M}_{\Sigma,m}^{q,\sigma}(\chi)$  and  $\mathcal{M}_{\Sigma,m}^{q,\sigma}(\psi)$  of the function class  $\Sigma_m$  and obtain estimates coefficient  $|\rho_{m+1}|$  and  $|\rho_{2m+1}|$  for functions in these two new subclasses.

**2. Main 4esults**

**Definition 4.** A function  $f(z)$  given by (3) is said to be in the class  $\mathcal{M}_{\Sigma,m}^{q,\sigma}(\chi)$  ( $m \in \mathcal{N}, 0 < q < 1, \sigma \geq 1, 0 < \chi \leq 1, (z, v) \in \Omega$ ) if

$$f \in \Sigma \quad \text{and} \quad |\arg(\mathcal{D}_q f(z))^\sigma| < \frac{\chi\pi}{2}, \tag{8}$$

and

$$|\arg(\mathcal{D}_q g(v))^\sigma| < \frac{\chi\pi}{2}, \tag{9}$$

where  $g(v)$  is given by (2).

**Remark 1.** We have the class  $\lim_{q \rightarrow 1^-} \mathcal{M}_{\Sigma,1}^\sigma(\chi) = \mathcal{M}_\Sigma^\sigma(\chi)$  which was introduced and studied by Girgaonkar et al., [29].

**Remark 2.** We have the class  $\lim_{q \rightarrow 1^-} \mathcal{M}_{\Sigma,1}^1(\chi) = \mathcal{M}_\Sigma(\chi)$  which was introduced and studied by Srivastava et al., [11].

**Theorem 1.** Let  $f(z) \in \mathcal{M}_{\Sigma,m}^{q,\sigma}(\chi)$ , ( $m \in \mathcal{N}, 0 < q < 1, \sigma \geq 1, 0 < \chi \leq 1, (z, v) \in \Omega$ ) be given (3). Then

$$|\rho_{m+1}| \leq \frac{2\chi}{\sqrt{(m+1)\sigma\chi[2m+1]_q - (\chi - \sigma)\sigma[m+1]_q^2}}, \tag{10}$$

and

$$|\rho_{2m+1}| \leq \frac{2\chi}{\sigma[2m+1]_q} + \frac{2(m+1)\chi^2}{\sigma^2[m+1]_q^2}. \tag{11}$$

**Proof.** Using inequalities (1) and (9), we get

$$(\mathcal{D}_q f(z))^\sigma = [\tau(z)]^\chi, \tag{12}$$

and

$$(\mathcal{D}_q g(v))^\sigma = [\zeta(v)]^\chi \tag{13}$$

respectively, where  $\tau(z)$  and  $\zeta(v)$  in  $\mathcal{P}$  are given by the following series

$$\tau(z) = 1 + \tau_m z^m + \tau_{2m} z^{2m} + \tau_{3m} z^{3m} + \dots, \tag{14}$$

and

$$\zeta(v) = 1 + \zeta_m v^m + \zeta_{2m} v^{2m} + \zeta_{3m} v^{3m} + \dots. \tag{15}$$

Clearly,

$$[\tau(z)]^\chi = 1 + \chi\tau_m z^m + \left(\chi\tau_{2m} + \frac{\chi(\chi-1)}{2}\tau_m^2\right) z^{2m} + \dots,$$

and

$$[\zeta(v)]^\chi = 1 + \chi\zeta_m v^m + \left(\chi\zeta_{2m} + \frac{\chi(\chi-1)}{2}\zeta_m^2\right) v^{2m} + \dots.$$

Also

$$(\mathcal{D}_q f(z))^\sigma = 1 + \sigma[m+1]_q \rho_{m+1} z^m + \left(\sigma[2m+1]_q \rho_{2m+1} + \frac{\sigma(\sigma-1)}{2}[m+1]_q^2 \rho_{m+1}^2\right) z^{2m} + \dots,$$

and

$$\begin{aligned}
 (\mathcal{D}_q g(v))^\sigma &= 1 - \sigma[m+1]_q \rho_{m+1} v^m - \sigma[2m+1]_q \rho_{2m+1} v^{2m} \\
 &\quad + \left( \sigma(m+1)[2m+1]_q \rho_{m+1}^2 + \frac{\sigma(\sigma-1)}{2} [m+1]_q^2 \rho_{m+1}^2 \right) v^{2m} + \dots
 \end{aligned}$$

Comparing the coefficients in (12) and (13), we have

$$\sigma[m+1]_q \rho_{m+1} = \chi \tau_m, \tag{16}$$

$$\sigma[2m+1]_q \rho_{2m+1} + \frac{\sigma(\sigma-1)}{2} [m+1]_q^2 \rho_{m+1}^2 = \chi \tau_{2m} + \frac{\chi(\chi-1)}{2} \tau_m^2, \tag{17}$$

$$-\sigma[m+1]_q \rho_{m+1} = \chi \zeta_m, \tag{18}$$

$$-\sigma[2m+1]_q \rho_{2m+1} + \left( \sigma(m+1)[2m+1]_q + \frac{\sigma(\sigma-1)}{2} [m+1]_q^2 \right) \rho_{m+1}^2 = \chi \zeta_{2m} + \frac{\chi(\chi-1)}{2} \zeta_m^2. \tag{19}$$

From (16) and (18), we obtain

$$\tau_m = -\zeta_m, \tag{20}$$

and

$$2\sigma[m+1]_q^2 \rho_{m+1}^2 = \chi^2 (\tau_m^2 + \zeta_m^2). \tag{21}$$

Further from (17), (19) and (21), we obtain that

$$\sigma(\sigma-1)\chi[m+1]_q^2 \rho_{m+1}^2 + (m+1)\sigma\chi[2m+1]_q \rho_{m+1}^2 - (\chi-1)\sigma^2[m+1]_q^2 \rho_{m+1}^2 = \chi^2 (\tau_{2m} + \zeta_{2m}).$$

Therefore, we have

$$\rho_{m+1}^2 = \frac{\chi^2 (\tau_{2m} + \zeta_{2m})}{\sigma[m+1]_q^2 (\sigma-\chi) + (m+1)\sigma\chi[2m+1]_q}. \tag{22}$$

By applying Lemma 1 for the coefficients  $\tau_{2m}$  and  $\zeta_{2m}$ , then we have

$$|\rho_{m+1}| \leq \frac{2\chi}{\sqrt{(m+1)\sigma\chi[2m+1]_q - (\chi-\sigma)\sigma[m+1]_q^2}}.$$

Also, to find the bound on  $|\rho_{2m+1}|$ , using the relation (19) and (17), we obtain

$$2\sigma[2m+1]_q \rho_{2m+1} - (m+1)\sigma[2m+1]_q \rho_{m+1}^2 = \chi(\tau_{2m} - \zeta_{2m}) + \frac{\chi(\chi-1)}{2} (\tau_m^2 - \zeta_m^2). \tag{23}$$

It follows from (20), (21) and (23),

$$\rho_{2m+1} = \frac{(m+1)\chi^2 \tau_m^2}{2\sigma^2[m+1]_q^2} + \frac{\chi(\tau_{2m} - \zeta_{2m})}{2\sigma[2m+1]_q}. \tag{24}$$

Applying Lemma 1 for the coefficients  $\tau_m, \tau_{2m}, \zeta_m, \zeta_{2m}$ , then we have

$$|\rho_{2m+1}| \leq \frac{2\chi}{\sigma[2m+1]_q} + \frac{2(m+1)\chi^2}{\sigma^2[m+1]_q^2}.$$

□

Choosing  $q \rightarrow 1^{-1}$  in Theorem 1, we get the following result:

**Corollary 1.** Let  $f(z) \in \mathcal{M}_{\Sigma, m}^\sigma(\chi)$ ,  $(m \in \mathcal{N}, \sigma \geq 1, 0 < \chi \leq 1, (z, v) \in \Omega)$  be given (3). Then

$$|\rho_{m+1}| \leq \frac{2\chi}{\sqrt{(m+1)[\sigma\chi m + \sigma^2 m + \sigma^2]}} \tag{25}$$

and

$$|\rho_{2m+1}| \leq \frac{2\chi}{\sigma(2m+1)} + \frac{2\chi^2}{\sigma^2(m+1)}. \tag{26}$$

Choosing  $m = 1$  (One-fold case) in Theorem 1, we get the following result:

**Corollary 2.** Let  $f(z) \in \mathcal{M}_{\Sigma}^{q,\sigma}(\chi)$ , ( $0 < q < 1, \sigma \geq 1, 0 < \chi \leq 1, (z, v) \in \Omega$ ) be given (1). Then

$$|\rho_2| \leq \frac{2\chi}{\sqrt{2\sigma\chi[3]_q - (\chi - \sigma)\sigma[2]_q^2}}, \tag{27}$$

and

$$|\rho_3| \leq \frac{2\chi}{\sigma[3]_q} + \frac{4\chi^2}{\sigma^2[2]_q^2}, \tag{28}$$

Choosing  $q \rightarrow 1^{-1}$  in Corollary 2, we get the following result:

**Corollary 3.** [29] Let  $f(z) \in \mathcal{M}_{\Sigma}^{\sigma}(\chi)$ , ( $\sigma \geq 1, 0 < \chi \leq 1, (z, v) \in \Omega$ ) be given (1). Then

$$|\rho_2| \leq \frac{2\chi}{\sqrt{2\sigma(2\sigma + \chi)}}, \tag{29}$$

and

$$|\rho_3| \leq \frac{\chi(2\sigma + 3\chi)}{3\sigma^2}. \tag{30}$$

**Remark 3.** For one-fold case, we have  $\lim_{q \rightarrow 1^{-1}} \mathcal{M}_{\Sigma,1}^{q,1}(\chi) = \mathcal{M}_{\Sigma}(\chi)$ , and we can get the results of Srivastava et al., [11].

**Definition 5.** A function  $f(z)$  given by (3) is said to be in the class  $\mathcal{M}_{\Sigma,m}^{q,\sigma}(\psi)$  ( $m \in \mathcal{N}, 0 < q < 1, \sigma \geq 1, 0 \leq \psi < 1, (z, v) \in \Omega$ ) if

$$f \in \Sigma \quad \text{and} \quad \Re[(\mathcal{D}_q f(z))^{\sigma}] > \psi, \tag{31}$$

and

$$\Re[(\mathcal{D}_q g(v))^{\sigma}] > \psi, \tag{32}$$

where  $g(v)$  is given by (2).

**Remark 4.** We have the class  $\lim_{q \rightarrow 1^{-1}} \mathcal{M}_{\Sigma,1}^{\sigma}(\psi) = \mathcal{M}_{\Sigma}^{\sigma}(\chi)$  which was introduced and studied by Girgaonkar et al., [29].

**Remark 5.** We have the class  $\lim_{q \rightarrow 1^{-1}} \mathcal{M}_{\Sigma,1}^1(\psi) = \mathcal{M}_{\Sigma}(\chi)$  which was introduced and studied by Srivastava et al., [11].

**Theorem 2.** Let  $f(z) \in \mathcal{M}_{\Sigma,m}^{q,\sigma}(\psi)$ , ( $m \in \mathcal{N}, 0 < q < 1, \sigma \geq 1, 0 \leq \psi < 1, (z, v) \in \Omega$ ) be given (3). Then

$$|\rho_{m+1}| \leq \min \left\{ \frac{2(1-\psi)}{\sigma[m+1]_q}, 2\sqrt{\frac{1-\psi}{\sigma(\sigma-1)[m+1]_q^2 + (m+1)\sigma[2m+1]_q}} \right\}, \tag{33}$$

and

$$|\rho_{2m+1}| \leq \frac{2(m+1)(1-\psi)}{\sigma(\sigma-1)[m+1]_q^2 + (m+1)\sigma[2m+1]_q} + \frac{2(1-\psi)}{\sigma[2m+1]_q}. \tag{34}$$

**Proof.** Using inequalities (31) and (32), we get

$$(\mathcal{D}_q f(z))^{\sigma} = \psi + (1-\psi)\tau(z), \tag{35}$$

and

$$(\mathcal{D}_q g(v))^\sigma = \psi + (1 - \psi)\zeta(v), \tag{36}$$

where  $\tau(z)$  and  $\zeta(v)$  in  $\mathcal{P}$  are given by the following series

$$\tau(z) = 1 + \tau_m z^m + \tau_{2m} z^{2m} + \tau_{3m} z^{3m} + \dots,$$

and

$$\zeta(v) = 1 + \zeta_m v^m + \zeta_{2m} v^{2m} + \zeta_{3m} v^{3m} + \dots.$$

Clearly,

$$\psi + (1 - \psi)\tau(z) = 1 + (1 - \psi)\tau_m z^m + (1 - \psi)\tau_{2m} z^{2m} + \dots,$$

and

$$\psi + (1 - \psi)\zeta(v) = 1 + (1 - \psi)\zeta_m v^m + (1 - \psi)\zeta_{2m} v^{2m} + \dots.$$

Also

$$(\mathcal{D}_q f(z))^\sigma = 1 + \sigma[m + 1]_q \rho_{m+1} z^m + \left( \sigma[2m + 1]_q \rho_{2m+1} + \frac{\sigma(\sigma - 1)}{2} [m + 1]_q^2 \rho_{m+1}^2 \right) z^{2m} + \dots,$$

and

$$\begin{aligned} (\mathcal{D}_q g(v))^\sigma &= 1 - \sigma[m + 1]_q \rho_{m+1} v^m - \sigma[2m + 1]_q \rho_{2m+1} v^{2m} \\ &+ \left( \sigma(m + 1)[2m + 1]_q \rho_{m+1}^2 + \frac{\sigma(\sigma - 1)}{2} [m + 1]_q^2 \rho_{m+1}^2 \right) v^{2m} + \dots. \end{aligned}$$

Now comparing the coefficients in (35) and (36), we get

$$\sigma[m + 1]_q \rho_{m+1} = (1 - \psi)\tau_m, \tag{37}$$

$$\sigma[2m + 1]_q \rho_{2m+1} + \frac{\sigma(\sigma - 1)}{2} [m + 1]_q^2 \rho_{m+1}^2 = (1 - \psi)\tau_{2m}, \tag{38}$$

$$-\sigma[m + 1]_q \rho_{m+1} = (1 - \psi)\zeta_m, \tag{39}$$

$$-\sigma[2m + 1]_q \rho_{2m+1} + \left( \sigma(m + 1)[2m + 1]_q + \frac{\sigma(\sigma - 1)}{2} [m + 1]_q^2 \right) \rho_{m+1}^2 = (1 - \psi)\zeta_{2m}. \tag{40}$$

From (37) and (39), we obtain

$$\tau_m = -\zeta_m, \tag{41}$$

and

$$2\sigma[m + 1]_q^2 \rho_{m+1}^2 = (1 - \psi)^2 (\tau_m^2 + \zeta_m^2). \tag{42}$$

Also, from (38) and (40), we get

$$\sigma(\sigma - 1)\chi[m + 1]_q^2 \rho_{m+1}^2 + (m + 1)\sigma[2m + 1]_q \rho_{m+1}^2 = (1 - \psi)(\tau_{2m} + \zeta_{2m}). \tag{43}$$

Applying the Lemma 1 for the coefficients  $\tau_m, \tau_{2m}, \zeta_m, \zeta_{2m}$ , we find that

$$|\rho_{m+1}| \leq 2\sqrt{\frac{(1 - \psi)}{\sigma(\sigma - 1)[m + 1]_q^2 + (m + 1)\sigma[2m + 1]_q}}.$$

Also, to find the bound on  $|\rho_{2m+1}|$ , using the relation (40) and (38), we obtain

$$-(m + 1)\sigma[2m + 1]_q \rho_{m+1}^2 + 2\sigma[2m + 1]_q \rho_{2m+1} = (1 - \psi)(\tau_{2m} - \zeta_{2m}), \tag{44}$$

or equivalently

$$\rho_{2m+1} = \frac{(1 - \psi)(\tau_{2m} - \zeta_{2m})}{2\sigma[2m + 1]_q} + \frac{(m + 1)}{2} \rho_{m+1}^2. \tag{45}$$

By substituting the value of  $\rho_{m+1}^2$  from (42), we have

$$\rho_{2m+1} = \frac{(1 - \psi)(\tau_{2m} - \zeta_{2m})}{2\sigma[2m + 1]_q} + \frac{(m + 1)(1 - \psi)^2(\tau_m^2 + \zeta_m^2)}{4\sigma^2[m + 1]_q^2}. \tag{46}$$

Applying the Lemma 1 for the coefficients  $\tau_m, \tau_{2m}, \zeta_m, \zeta_{2m}$ , we get

$$|\rho_{2m+1}| \leq \frac{2(1 - \psi)}{\sigma[2m + 1]_q} + \frac{2(m + 1)(1 - \psi)^2}{2\sigma^2[m + 1]_q^2}.$$

Also, by using (43) and (45), and applying Lemma 1 we obtain

$$|\rho_{2m+1}| \leq \frac{2(m + 1)(1 - \psi)}{\sigma(\sigma - 1)[m + 1]_q^2 + (m + 1)\sigma[2m + 1]_q} + \frac{2(1 - \psi)}{\sigma[2m + 1]_q}.$$

This complete the proof.  $\square$

Choosing  $q \rightarrow 1^{-1}$  in Theorem 2, we get the following result:

**Corollary 4.** Let  $f(z) \in \mathcal{M}_{\Sigma, m}^\sigma(\psi)$ ,  $(m \in \mathcal{N}, \sigma \geq 1, 0 \leq \psi < 1, (z, v) \in \Omega)$  be given (3). Then

$$|\rho_{m+1}| \leq \begin{cases} 2\sqrt{\frac{(1-\psi)}{\sigma(\sigma-1)[m+1]^2+(m+1)\sigma[2m+1]}} & 0 \leq \psi \leq \frac{m}{1+2m}, \\ \frac{2(1-\psi)}{\sigma[m+1]} & \frac{m}{1+2m} \leq \psi < 1, \end{cases}$$

and

$$|\rho_{2m+1}| \leq \frac{2(m + 1)(1 - \psi)}{\sigma(\sigma - 1)[m + 1]^2 + (m + 1)\sigma[2m + 1]} + \frac{2(1 - \psi)}{\sigma[2m + 1]}.$$

For one fold case, Corollary 4, yields the following Corollary:

**Corollary 5.** Let  $f(z) \in \mathcal{M}_\Sigma^\sigma(\psi)$ ,  $(\sigma \geq 1, 0 \leq \psi < 1, (z, v) \in \Omega)$  be given (1). Then

$$|\rho_2| \leq \begin{cases} \sqrt{\frac{2(1-\psi)}{\sigma(2\sigma+1)}} & 0 \leq \psi \leq \frac{1}{3}, \\ \frac{(1-\psi)}{\sigma} & \frac{1}{3} \leq \psi < 1, \end{cases}$$

and

$$|\rho_3| \leq \frac{(1 - \psi)(2\sigma - 3\psi + 3)}{3\sigma^2}.$$

**Remark 6.** Corollary 5 gives above is the improvement of the estimates for coefficients on  $|\rho_2|$  and  $|\rho_3|$  investigated by Girgaonkar *et al.*, [29].

**Corollary 6.** [29] Let  $f(z) \in \mathcal{M}_\Sigma^\sigma(\psi)$ ,  $(\sigma \geq 1, 0 \leq \psi < 1, (z, v) \in \Omega)$  be given (1). Then

$$|\rho_2| \leq \sqrt{\frac{2(1 - \psi)}{\sigma(2\sigma + 1)'}}$$

and

$$|\rho_3| \leq \frac{(1 - \psi)(2\sigma - 3\psi + 3)}{3\sigma^2}.$$

Taking  $\sigma = 1$  in Corollary 7, we get the following result:



**Corollary 7.** [11] Let  $f(z) \in \mathcal{M}_{\Sigma}^{\sigma}(\psi)$ , ( $\sigma \geq 1, 0 \leq \psi < 1, (z, v) \in \Omega$ ) be given (1). Then

$$|\rho_2| \leq \sqrt{\frac{2(1-\psi)}{3}},$$

and

$$|\rho_3| \leq \frac{(1-\psi)(5-3\psi)}{3}.$$

### 3. Conclusion

In this present paper, two new subclasses indicated by  $\mathcal{M}_{\Sigma, m}^{q, \sigma}(\chi)$  and  $\mathcal{M}_{\Sigma, m}^{q, \sigma}(\psi)$  of function class of  $\mathcal{E}_m$  was obtained and worked on. Also, the estimates coefficients for  $|\rho_{m+1}|$  and  $|\rho_{2m+1}|$  of functions in these classes are determined.

**Conflicts of Interest:** "The author declares no conflict of interest."

### References

- [1] Duren, P. L. (2001). *Univalent Functions* (Vol. 259). Springer, New York, NY, USA.
- [2] Brannan, D. A., & Taha, T. S. (1988). On some classes of bi-univalent functions. In *Mathematical Analysis and Its Applications* (pp. 53-60). Pergamon.
- [3] Frasin, B. A., & Aouf, M. K. (2011). New subclasses of bi-univalent functions. *Applied Mathematics Letters*, 24(9), 1569-1573.
- [4] Lewin, M. (1967). On a coefficient problem for bi-univalent functions. *Proceedings of the American Mathematical Society*, 18(1), 63-68.
- [5] Sakar, F. M., & Güney, M. O. (2018). Coefficient estimates for certain subclasses of  $m$ -fold symmetric bi-univalent functions defined by the  $q$ -derivative operator. *Konuralp Journal of Mathematics*, 6(2), 279-285.
- [6] Patil, A. B., & Naik, U. H. (2018). Bounds on initial coefficients for a new subclass of bi-univalent functions. *New Trends in Mathematical Sciences*, 6(1), 85-90.
- [7] Shaba, T.G., Ibrahim, A. A., & Jimoh, A. A. (2020). On a new subclass of bi-pseudo-starlike functions defined by Frasin differential operator. *Advances in Mathematics: Scientific Journal*, 9(7) (2020), 4829-4841.
- [8] Shaba, T.G. (2020). On some new subclass of bi-univalent functions associated with the Opoola differential operator. *Open Journal of Mathematical Analysis*, 4(2), 74-79.
- [9] Shaba, T. G. (2020). Subclass of bi-univalent functions satisfying subordinate conditions defined by Frasin differential operator. *Turkish Journal of Inequalities*, 4(2), 50-58.
- [10] Shaba, T. G. (2020). Certain new subclasses of analytic and bi-univalent functions using Salagean operator. *Asia Pacific Journal of Mathematics*, 7(29), 7-29.
- [11] Srivastava, H. M., Mishra, A. K., & Gochhayat, P. (2010). Certain subclasses of analytic and bi-univalent functions. *Applied mathematics letters*, 23(10), 1188-1192.
- [12] Srivastava, H. M., Sivasubramanian, S., & Sivakumar, R. (2014). Initial coefficient bounds for a subclass of  $m$ -fold symmetric bi-univalent functions. *Tbilisi Mathematical Journal*, 7(2), 1-10.
- [13] Akgül, A., & Campus, U. (2017). On the coefficient estimates of analytic and bi-univalent  $m$ -fold symmetric functions. *Mathematica Aeterna*, 7(3), 253-260.
- [14] Altinkaya, S., & Yalçın, S. (2018). On some subclasses of  $m$ -fold symmetric bi-univalent functions. *Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics*, 67(1), 29-36.
- [15] Bulut, S. (2016). Coefficient estimates for general subclasses of  $m$ -fold symmetric analytic bi-univalent functions. *Turkish Journal of Mathematics*, 40(6), 1386-1397.
- [16] Hamidi, S. G., & Jahangiri, J. M. (2014). Unpredictability of the coefficients of  $m$ -fold symmetric bi-starlike functions. *International Journal of Mathematics*, 25(07), 1450064.
- [17] Eker, S. S. (2016). Coefficient bounds for subclasses of  $m$ -fold symmetric bi-univalent functions. *Turkish Journal of Mathematics*, 40(3), 641-646.
- [18] Jackson, D. O., Fukuda, T., Dunn, O., & Majors, E. (1910). On  $q$ -definite integrals. *Quarterly Journal of Pure and Applied Mathematics*, 41, 193-203.
- [19] Jackson, F. H. (1909). XI. on  $q$ -functions and a certain difference operator. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, 46(2), 253-281.
- [20] Akgül, A. (2018, January). Finding initial coefficients for a class of bi-univalent functions given by  $q$ -derivative. In *AIP Conference Proceedings* (Vol. 1926, No. 1, p. 020001). AIP Publishing LLC.

- [21] Aldweby, H., & Darus, M. (2013). A subclass of harmonic  $\mathfrak{U}$  univalent functions associated with  $q$ -analogue of Dziok-Srivastava operator. *ISRN Mathematical Analysis, 2013*, Article ID 382312.
- [22] Aldweby, H., & Darus, M. (2017). Coefficient estimates for initial Taylor-Maclaurin coefficients for a subclass of analytic and bi-univalent functions associated with  $q$ -derivative operator. *Recent Trends in Pure and Applied Mathematics, 2017*, 109-117
- [23] Aral, A., Gupta, V., & Agarwal, R. P. (2013). *Applications of  $q$ -calculus in operator theory* (p. 262). New York: Springer.
- [24] Bulut, S. (2017). Certain subclasses of analytic and bi-univalent functions involving the  $q$ -derivative operator. *Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics, 66(1)*, 108-114.
- [25] Mohammed, A., & Darus, M. (2013). A generalized operator involving the  $q$ -hypergeometric function. *Matematiski Vesnik, 65(254)*, 454-465.
- [26] Seoudy, T. M., & Aouf, M. K. (2014). Convolution properties for  $\mathfrak{C}$  certain classes of analytic functions defined by  $q$ -derivative operator, *Abstract and Applied Analysis, 2014*, Article ID 846719.
- [27] Seoudy, T. M., & Aouf, M. K. (2016). Coefficient estimates of new classes of  $q$ -starlike and  $q$ -convex functions of complex order. *Journal of Mathematical Inequalities, 10(1)*, 135-145.
- [28] Babalola, K.O.(2013). On  $\lambda$ -pseudo-starlike function. *Journal of Classical Analysis, 3*, 137-147.
- [29] Girgaonkar, V. B., Joshi, S. B., & Yadav, P. P. (2017). Certain special subclasses of analytic function associated with bi-univalent functions. *Palestine Journal of Mathematics, 6(2)*, 617-623.



© 2021 by the authors; licensee PSRP, Lahore, Pakistan. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).