

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/356524015>

Coefficient functionals for a class of bounded turning functions related to modified sigmoid function

Article in *AIMS Mathematics* · November 2021

DOI: 10.3934/math.2022173

CITATIONS

3

READS

297

5 authors, including:



Muhammad Ghaffar Khan

Kohat University of Science and Technology

46 PUBLICATIONS 366 CITATIONS

[SEE PROFILE](#)



Timilehin Gideon Shaba

Landmark University

57 PUBLICATIONS 369 CITATIONS

[SEE PROFILE](#)



Bakhtiar Ahmad

Govt. Degree College Mardan

44 PUBLICATIONS 333 CITATIONS

[SEE PROFILE](#)



Wali Khan Mashwani

Kohat University of Science and Technology

184 PUBLICATIONS 1,792 CITATIONS

[SEE PROFILE](#)



Research article

Coefficient functionals for a class of bounded turning functions related to modified sigmoid function

Muhammad Ghaffar Khan¹, Nak Eun Cho^{2,*}, Timilehin Gideon Shaba³, Bakhtiar Ahmad⁴ and Wali Khan Mashwani¹

¹ Institute of Numerical Sciences, Kohat university of science and technology, Kohat, Pakistan

² Department of Applied Mathematics, Pukyong National University Busan 48513, Korea

³ Department of Mathematics, University of Ilorin, P. M. B. 1515, Ilorin, Nigeria

⁴ Govt. Degree College Mardan, Mardan 23200, Pakistan

* **Correspondence:** Email: necho@pknu.ac.kr.

Abstract: The main objective of the present article is to define the class of bounded turning functions associated with modified sigmoid function. Also we investigate and determine sharp results for the estimates of four initial coefficients, Fekete-Szegő functional, the second-order Hankel determinant, Zalcman conjecture and Krushkal inequality. Furthermore, we evaluate bounds of the third and fourth-order Hankel determinants for the class and for the 2-fold and 3-fold symmetric functions.

Keywords: analytic functions; modified sigmoid function; subordination; Hankel determinant; 2-fold and 3-fold symmetric function; Zalcman conjecture; Krushkal inequality

Mathematics Subject Classification: 30C45, 30D30

1. Introduction

Let \mathcal{A} represent the collections of analytic functions defined in open unit disc $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ whose normalization is of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (z \in \mathbb{D}). \tag{1.1}$$

Let \mathcal{S} denote the subclass of \mathcal{A} comprising of functions of the form (1.1) which are also univalent in \mathbb{D} .

Let \mathcal{P} represent the class of all functions p that are analytic in \mathbb{D} with $\Re(p(z)) > 0$ and has the series representation

$$p(z) = 1 + \sum_{n=1}^{\infty} c_n z^n \quad (z \in \mathbb{D}). \tag{1.2}$$

Next we recall the definition of subordination. For two functions $h_1, h_2 \in \mathcal{A}$, we say that h_1 is subordinate to h_2 and is symbolically written as $h_1 < h_2$ if there exists an analytic function w with the property $|w(z)| \leq |z|$ and $w(0) = 0$ such that $h_1(z) = h_2(w(z))$ for $z \in \mathbb{D}$. Further, if $h_2 \in \mathcal{S}$, then the condition becomes

$$h_1 < h_2 \Leftrightarrow h_1(0) = h_2(0) \text{ and } h_1(\mathbb{D}) \subset h_2(\mathbb{D}).$$

Now we consider the following class $\mathcal{S}^*(\varphi)$ as follows:

$$\mathcal{S}^*(\varphi) = \left\{ f \in \mathcal{A} : \frac{zf'(z)}{f(z)} < \varphi(z) \right\}, \quad (1.3)$$

where φ is an analytic univalent function with positive real part in \mathbb{D} , $\varphi(\mathbb{U})$ is symmetric about the real axis and starlike with respect to $\varphi(0) = 1$ and $\varphi'(0) > 0$. The class $\mathcal{S}^*(\varphi)$ was introduced by Ma and Minda [20]. In particular, if we take $\varphi(z) = (1+z)/(1-z)$, then the class $\mathcal{S}^*(\varphi)$ is the well-known class of starlike functions. If we vary the function φ on the right hand side of (1.3), then we obtain some several subclasses of \mathcal{S} whose image domains have some interesting geometrical configurations as follows:

- (1) The class $\mathcal{S}^*(\varphi)$ with $\varphi(z) = 1 + \sin z$ is introduced and studied by Cho et al. [6].
- (2) The class $\mathcal{S}^*(\varphi)$ with $\varphi(z) = 1 + z - \frac{1}{3}z^3$, which is a nephroid shaped domain, was introduced and investigated by Wani and Swaminathan [39].
- (3) The class $\mathcal{S}^*(\varphi)$ with $\varphi(z) = \sqrt{1+z}$, which is bounded by lemniscate of Bernoulli in right half plan, was developed by Sokól and Stankiewicz [30].
- (4) The class $\mathcal{S}^*(\varphi)$ with $\varphi(z) = 1 + \frac{4}{3}z + \frac{2}{3}z^2$ was introduced by Sharma et al. [29].
- (5) The class $\mathcal{S}^*(\varphi)$ with $\varphi(z) = e^z$ was introduced and studied by Mendiratta et al. [21].
- (6) The class $\mathcal{S}^*(\varphi)$ with $\varphi(z) = z + \sqrt{1+z^2}$, which maps \mathbb{D} to crescent shaped region, was introduced by Raina and Sokól [26].

Also we note that lately many subclasses of starlike functions are introduced see [7, 9, 12] by choosing some particular functions such as functions associated with Bell numbers, shell-like curve connected with Fibonacci numbers, functions connected with conic domains and rational functions instead of φ in (1.3).

Pommerenke [24, 25] introduced the Hankel determinant $H_{q,n}(f)$ for function $f \in \mathcal{S}$ of the form (1.1), where the parameters $q, n \in \mathbb{N} = \{1, 2, 3, \dots\}$ as follows:

$$H_{q,n}(f) = \begin{vmatrix} a_n & a_{n+1} & \dots & a_{n+q-1} \\ a_{n+1} & a_{n+2} & \dots & a_{n+q} \\ \vdots & \vdots & \dots & \vdots \\ a_{n+q-1} & a_{n+q} & \dots & a_{n+2q-2} \end{vmatrix}. \quad (1.4)$$

The Hankel determinants for different orders are obtained for different values of q and n . When $q = 2$ and $n = 1$, the determinant is

$$|H_{2,1}(f)| = \begin{vmatrix} a_1 & a_2 \\ a_2 & a_3 \end{vmatrix} = |a_3 - a_2^2|, \text{ where } a_1 = 1.$$

Note that $H_{2,1}(f) = a_3 - a_2^2$, is the classical Fekete-Szegő functional. For various subclasses of \mathcal{A} , the best possible value of the upper bound for $|H_{2,1}(f)|$ was investigated by different authors (see [13–15] for details). Furthermore, when $q = 2$ and $n = 2$, the second Hankel determinant is

$$H_{2,2}(f) = \begin{vmatrix} a_2 & a_3 \\ a_3 & a_4 \end{vmatrix} = a_2a_4 - a_3^2.$$

The upper bound of $|H_{2,2}(f)|$ has been studied by several authors in the last few decades. For instance, the readers may refer to the works of Hayman [11], the Noonan and Thomas [22], Ohran et al. [23] and Shi et al. [34]. Moreover, Babalola [3] studied the Hankel determinant $H_{3,1}(f)$ for some subclasses of analytic functions. For some recent works on third order Hankel determinant we may refer the interested reader to such more recent works as (for example) [28, 32, 38]. The bound of the fourth Hankel determinant for a class of analytic functions with bounded turning associated with cardioid domain was approximated by Srivastava et al. in [37]. It should be remarked that a wide variety of applications of Hankel systems arise in linear filtering theory, discrete inverse scattering, and discretization of certain integral equations arising in mathematical physics [40].

Evaluating these Hankel determinants for various new subclasses has been an attracting area lately. One such field of interest is the Quantum Calculus (q -calculus), which is a generalization of classical calculus by replacing the limit by a parameter q . For the basics and preliminaries, the readers are advised to see the works and expositions in [31, 35, 37]. It is important to mention here the work on a q -differential operator by Srivastava et al. [33], in which they determined the upper bound of second Hankel determinant for a subclass of bi-univalent functions in q -analogue. Recently, the upper bound estimate for q -analogue of a subclass of starlike functions in connection with exponential function were evaluated in [36].

Recently, a class of starlike functions associated with Modified sigmoid function was defined by Goel and Kumar [10], i.e,

$$\mathcal{S}_{S_G}^* = \left\{ f \in \mathcal{S} : \frac{zf'(z)}{f(z)} < \frac{2}{1+e^{-z}} \right\} \quad (z \in \mathbb{D}).$$

Motivated by all the works mentioned above and [4], in this article we introduce and investigate the class \mathcal{R}_{S_G} , which is defined as follows:

$$\mathcal{R}_{S_G} = \left\{ f \in \mathcal{S} : f'(z) < \frac{2}{1+e^{-z}} \right\} \quad (z \in \mathbb{D}). \quad (1.5)$$

We also establish some sharp results such as coefficient bounds, Fekete-Szegő inequality, second-order determinant, Zalcman conjecture and Krushkal inequality for functions belonging to the class \mathcal{R}_{S_G} . Moreover, we estimate bounds of the third and forth-order Hankel determinants for this class \mathcal{R}_{S_G} and for the 2-fold and 3-fold symmetric functions.

2. A set of lemmas

For the proofs of our main findings, we need the following lemmas.

Lemma 1. Let $p \in \mathcal{P}$ have the series expansion of the form (1.2). Then, for x and σ with $|x| \leq 1, |\sigma| \leq 1$, such that

$$2c_2 = c_1^2 + x(4 - c_1^2), \quad (2.1)$$

$$4c_3 = c_1^3 + 2(4 - c_1^2)lc_1x - c_1(4 - c_1^2)x^2 + 2(4 - c_1^2)(1 - |x|^2)\sigma. \quad (2.2)$$

We note that (2.1) and (2.2) are taken from [18].

Lemma 2. If $p \in \mathcal{P}$ and has the series of the form (1.2), then

$$|c_{n+k} - \mu c_n c_k| \leq 2, \quad 0 \leq \mu \leq 1, \quad (2.3)$$

$$|c_n| \leq 2 \text{ for } n \geq 1, \quad (2.4)$$

$$|c_2 - \zeta c_1^2| \leq 2 \max\{1, |2\zeta - 1|\}, \quad \zeta \in \mathbb{C}. \quad (2.5)$$

We note that the inequalities (2.3), (2.4) and (2.6) in the above can be found in [2, 25] and (2.5) is given by [13].

Lemma 3. [2] If $p \in \mathcal{P}$ and has the series of the form (1.2), then

$$|Jc_1^3 - Kc_1c_2 + Lc_3| \leq 2|J| + 2|K - 2J| + 2|J - K + L|, \quad (2.6)$$

where J, K and L are real numbers.

Lemma 4. [27] Let m, n, l and r satisfy the inequalities $0 < m < 1, 0 < r < 1$ and

$$8r(1-r) \left[(mn - 2l)^2 + (m(r+m) - n)^2 \right] + m(1-m)(n - 2rm)^2 \leq 4m^2(1-m)^2 r(1-r).$$

If $p \in \mathcal{P}$ and has power series (1.2), then

$$\left| lc_1^4 + rc_2^2 + 2mc_1c_3 - \frac{3}{2}nc_1^2c_2 - c_4 \right| \leq 2.$$

3. Bounds of $|H_{3,1}(f)|$ for the class \mathcal{R}_{S_G}

Theorem 1. Let $f \in \mathcal{R}_{S_G}$ and be of the form (1.1). Then

$$|a_2| \leq \frac{1}{4}, \quad (3.1)$$

$$|a_3| \leq \frac{1}{6}, \quad (3.2)$$

$$|a_4| \leq \frac{1}{8}, \quad (3.3)$$

$$|a_5| \leq \frac{1}{10}, \quad (3.4)$$

$$|a_6| \leq \frac{355}{288}, \quad (3.5)$$

$$|a_7| \leq \frac{381377}{2820}. \quad (3.6)$$

The first four inequalities are sharp for the functions defined below respectively

$$f_n(z) = \int_0^z \left(\frac{2}{1 + e^{-t}} \right) dt = z + \frac{1}{2(n+1)} z^{n+1} + \dots, \quad \text{where } n = 1, 2, 3, 4. \quad (3.7)$$

Proof. Let $f \in \mathcal{R}_{S_G}$. Then, (1.5) can be put in the form of Schwarz function $w(z)$ as

$$f'(z) = \frac{2}{1 + e^{-w(z)}} \quad (z \in \mathbb{D}). \quad (3.8)$$

Also, if $p \in \mathcal{P}$, then it may be written in terms of the Schwarz function w as

$$p(z) = 1 + c_1 z + c_2 z^2 + c_3 z^3 \dots = \frac{1 + w(z)}{1 - w(z)},$$

or equivalently,

$$w(z) = \frac{p(z) - 1}{p(z) + 1} = \frac{1}{2} c_1 z + \left(\frac{1}{2} c_2 - \frac{1}{4} c_1^2 \right) z^2 + \left(\frac{1}{8} c_1^3 - \frac{1}{2} c_2 c_1 + \frac{1}{2} c_3 \right) z^3 + \dots. \quad (3.9)$$

Now

$$f'(z) = 1 + 2a_2 z + 3a_3 z^2 + 4a_4 z^3 + 5a_5 z^4 + \dots, \quad (3.10)$$

By a simplification and using the series expansion (3.9), we have

$$\begin{aligned} \frac{2}{1 + e^{-w(z)}} &= 1 + \frac{c_1}{4} z + \left(\frac{c_2}{4} - \frac{c_1^2}{8} \right) z^2 + \left(\frac{11c_1^3}{192} - \frac{c_2 c_1}{4} + \frac{c_3}{4} \right) z^3 \\ &+ \left(-\frac{3}{128} c_1^4 + \frac{11}{64} c_1^2 c_2 - \frac{1}{4} c_3 c_1 - \frac{1}{8} c_2^2 + \frac{1}{4} c_4 \right) z^4 + \dots. \end{aligned} \quad (3.11)$$

Comparing (3.10) and (3.11), we get

$$a_2 = \frac{1}{8} c_1, \quad (3.12)$$

$$a_3 = \frac{1}{3} \left(\frac{1}{4} c_2 - \frac{1}{8} c_1^2 \right), \quad (3.13)$$

$$a_4 = \frac{1}{4} \left(\frac{11}{192} c_1^3 - \frac{1}{4} c_1 c_2 + \frac{1}{4} c_3 \right) \quad (3.14)$$

$$a_5 = -\frac{1}{20} \left(\frac{3}{32} c_1^4 - \frac{11}{16} c_1^2 c_2 + c_3 c_1 + \frac{1}{2} c_2^2 - c_4 \right). \quad (3.15)$$

$$a_6 = \frac{1}{18432} \left(\begin{array}{l} -5c_1^6 + 122c_1^4 c_2 - 288c_1^3 c_3 - 432c_1^2 c_2^2 + 528c_4 c_1^2 + 1056c_1 c_2 c_3 \\ -768c_5 c_1 + 176c_2^3 - 768c_4 c_2 - 384c_3^2 + 768c_6 \end{array} \right) \quad (3.16)$$

and

$$a_7 = \frac{1}{36\,126\,720} \left(\begin{array}{l} -2537c_1^7 - 50\,400c_1^5 c_2 + 204\,960c_1^4 c_3 + 409\,920c_1^3 c_2^2 - 483\,840c_4 c_1^3 \\ -1451\,520c_1^2 c_2 c_3 + 887\,040c_5 c_1^2 - 483\,840c_1 c_2^3 + 1774\,080c_4 c_1 c_2 + 887\,040c_1 c_3^2 \\ -1290\,240c_6 c_1 + 887\,040c_2^2 c_3 - 1290\,240c_5 c_2 - 1290\,240c_4 c_3 + 1290\,240c_7 \end{array} \right). \quad (3.17)$$

For a_2 , putting (2.4) in (3.12), we have

$$|a_2| \leq \frac{1}{4}.$$

For a_3 , simplifying (3.13), we get

$$a_3 = \frac{1}{12} \left(c_2 - \frac{c_1^2}{2} \right)$$

and applying (2.3), we have

$$|a_3| \leq \frac{1}{6}.$$

For a_4 , using (3.14), we obtain

$$|a_4| = \frac{1}{4} \left| \frac{11}{192} c_1^3 - \frac{1}{4} c_1 c_2 + \frac{1}{4} c_3 \right|. \quad (3.18)$$

By applying Lemma 3 to (3.18), we get

$$|a_4| \leq \frac{1}{4} \left[2 \left| \frac{11}{192} \right| + 2 \left| \frac{1}{4} - 2 \left(\frac{11}{192} \right) \right| + \left| \frac{11}{192} - \frac{1}{4} + \frac{1}{4} \right| \right] = \frac{1}{8}.$$

For a_5 , applying Lemma 4 to (3.15), we get

$$|a_5| \leq \frac{1}{10}.$$

For a_6 , re-arranging (3.16) and applying the triangle inequality, we get

$$|a_6| \leq \frac{1}{18432} \left[122 |c_1|^4 \left| c_2 - \frac{5}{122} c_1^2 \right| + 1056 |c_1| |c_3| \left| c_2 - \frac{3}{11} c_1^2 \right| + 528 |c_1|^2 \left| c_4 - \frac{9}{11} c_2^2 \right| \right. \\ \left. + 768 |c_6 - c_1 c_5| + 768 |c_2| \left| c_4 - \frac{88}{89} c_2^2 \right| + 384 |c_3|^2 \right].$$

By applying (2.3) and (2.4) to the above, we get

$$|a_6| \leq \frac{355}{288}.$$

For a_7 , re-arranging (3.17) and applying the triangle inequality, we get

$$|a_7| \leq \frac{1}{36\,126\,720} \left[204960 |c_1|^4 \left| c_3 - \frac{105}{427} c_1 c_2 \right| + 483840 |c_1|^3 \left| c_4 - \frac{61}{72} c_1^2 \right| + 1290240 |c_1| \left| c_6 - \frac{11}{16} c_1 c_5 \right| \right. \\ \left. + 1774080 |c_1| |c_2| \left| c_4 - \frac{9}{11} c_1 c_3 \right| + 887040 |c_2|^2 \left| c_3 - \frac{6}{11} c_1 c_2 \right| + 1290240 |c_7 - c_2 c_5| \right. \\ \left. + 1290240 |c_3| \left| c_4 - \frac{11}{16} c_1 c_3 \right| + 2537 |c_1|^7 \right].$$

Also by using (2.3) and (2.4) to the above, we obtain

$$|a_7| \leq \frac{381\,377}{282\,240}.$$

□

Next, we consider the Fekete-Szegő problem and the Hankel determinants for the class \mathcal{R}_{S_G} .

Theorem 2. If f of the form (1.1) belongs to \mathcal{R}_{S_G} , then

$$|a_3 - \zeta a_2^2| \leq \frac{1}{6} \max \left\{ 1, \frac{3|\zeta|}{8} \right\} \quad (\zeta \in \mathbb{C}). \quad (3.19)$$

The result is sharp for the function f_2 defined by (3.7) for $|\zeta| \leq 8/3$ and the function f_1 defined by (3.7) for $|\zeta| \geq 8/3$.

Proof. Using (3.12) and (3.13), we can write

$$|a_3 - \zeta a_2^2| = \left| \frac{c_2}{12} - \frac{c_1^2}{24} - \zeta \frac{c_1^2}{64} \right|.$$

By rearranging we have

$$|a_3 - \zeta a_2^2| = \frac{1}{12} \left| c_2 - \left(\frac{3\zeta + 8}{16} \right) c_1^2 \right|.$$

Applying (2.5) we get

$$|a_3 - \zeta a_2^2| \leq \frac{1}{12} \max \left\{ 2, 2 \left| 2 \left(\frac{3\zeta + 8}{16} \right) - 1 \right| \right\}.$$

Then with simple calculations, we obtain

$$|a_3 - \zeta a_2^2| \leq \frac{1}{6} \max \left\{ 1, \frac{3|\zeta|}{8} \right\}.$$

For the sharpness consider the function

$$f_2(z) = z + \frac{1}{6}z^3 - \frac{1}{168}z^7 + \dots, \quad (3.20)$$

which gives equality in (3.19) when $|\zeta| \leq \frac{8}{3}$, namely

$$|a_3 - \zeta a_2^2| = |a_3| = \frac{1}{6} = \frac{1}{6} \max \left\{ 1, \frac{3|\zeta|}{8} \right\}.$$

For the case $|\zeta| \geq \frac{8}{3}$ consider

$$f_1(z) = z + \frac{1}{4}z^2 - \frac{1}{96}z^4 + \dots,$$

which gives

$$|a_3 - \zeta a_2^2| = |\zeta a_2^2| = \frac{|\zeta|}{16}.$$

□

If we put $\zeta = 1$, then the above result becomes:

Corollary 1. If f of the form (1.1) belongs to \mathcal{R}_{S_G} , then

$$|a_3 - a_2^2| \leq \frac{1}{6}. \quad (3.21)$$

Theorem 3. If f of the form (1.1) belongs to \mathcal{R}_{S_G} , then

$$|a_2a_3 - a_4| \leq \frac{1}{8}. \quad (3.22)$$

The result is sharp for the function f_3 defined by (3.7).

Proof. From (3.12)–(3.14), we get

$$|a_2a_3 - a_4| = \frac{1}{16} \left| \frac{5}{16}c_1^3 - \frac{7}{6}c_2c_1 + c_3 \right|.$$

Using Lemma 3, we get the required result. \square

Theorem 4. If f of the form (1.1) belongs to \mathcal{R}_{S_G} , then

$$|H_{2,2}(f)| = |a_2a_4 - a_3^2| \leq \frac{1}{36}. \quad (3.23)$$

The result is sharp for the function f_2 defined by (3.7).

Proof. From (3.12)–(3.14), we have

$$H_{2,2}(f) = \frac{1}{18432}c_1^4 - \frac{1}{1152}c_1^2c_2 + \frac{1}{128}c_1c_3 - \frac{1}{144}c_2^2.$$

Applying (2.1) and (2.2) to express c_2 and c_3 in terms of $c_1 = c$, with $0 \leq c \leq 2$, we get

$$H_{2,2}(f) = -\frac{1}{6144}c^4 - \frac{1}{512}c^2(4 - c^2)x^2 - \frac{1}{576}(4 - c^2)^2x^2 + \frac{1}{256}c(4 - c^2)(1 - |x|^2)\sigma.$$

With the aid of the triangle inequality and replacing $|\sigma| \leq 1$, $|x| = b$, with $b \leq 1$, we obtain

$$|H_{2,2}(f)| \leq \frac{1}{6144}c^4 + \frac{1}{512}c^2(4 - c^2)b^2 + \frac{1}{576}(4 - c^2)^2b^2 + \frac{1}{256}c(4 - c^2)(1 - b^2) := \phi(c, b).$$

It is a simple calculation to show that $\frac{\partial\phi(c,b)}{\partial b} \geq 0$ on $[0, 1]$, so that $\phi(c, b) \leq \phi(c, 1)$. Putting $b = 1$ gives

$$|H_{2,2}(f)| \leq \frac{1}{6144}c^4 + \frac{1}{512}c^2(4 - c^2) + \frac{1}{576}(4 - c^2)^2 := \phi(c, 1).$$

Also, $\phi'(c, 1) = 0$, has only root $c = 0 \in [0, 2]$ and so $\phi''(0, 1) < 0$. Thus, the maximum value at $c = 0$ is

$$|H_{2,2}(f)| \leq \frac{1}{36}. \quad \square$$

Theorem 5. If $f \in \mathcal{A}$ belongs to \mathcal{R}_{S_G} , then

$$|a_2a_5 - a_3a_4| \leq \frac{217}{2880}. \quad (3.24)$$

Proof. From (3.12)–(3.15), we have

$$|a_2a_5 - a_3a_4| = \left| \frac{1}{92\,160}c_1^5 + \frac{23}{46\,080}c_1^3c_2 - \frac{7}{1920}c_3c_1^2 + \frac{1}{480}c_1c_2^2 + \frac{1}{160}c_4c_1 - \frac{1}{192}c_3c_2 \right|.$$

Rearranging the above term, we get

$$\begin{aligned} |a_2a_5 - a_3a_4| &= \left| \frac{1}{92\,160}c_1^5 - \frac{7}{1920}c_1^2 \left(c_3 - \frac{23}{168}c_1c_2 \right) - \frac{1}{192}c_2 \left(c_3 - \frac{2}{5}c_1c_2 \right) + \frac{1}{160}c_1c_4 \right| \\ &\leq \frac{1}{92\,160}|c_1|^5 + \frac{7}{1920}|c_1|^2 \left| c_3 - \frac{23}{168}c_1c_2 \right| + \frac{1}{192}|c_2| \left| c_3 - \frac{2}{5}c_1c_2 \right| + \frac{1}{160}|c_1||c_4|. \end{aligned}$$

Using (2.3) and (2.4), we get the required result. \square

Theorem 6. If $f \in \mathcal{A}$ belongs to \mathcal{R}_{S_G} , then

$$|a_5 - a_2a_4| \leq \frac{1}{10}. \quad (3.25)$$

The result is sharp for function f_4 defined by (3.7).

Proof. From (3.12)–(3.15), we have

$$|a_5 - a_2a_4| = \frac{1}{20} \left| \frac{199}{1536}c_1^4 - \frac{27}{32}c_1^2c_2 + \frac{37}{32}c_3c_1 + \frac{1}{2}c_2^2 - c_4 \right|.$$

By applying of Lemma 4, we get the desired result. \square

Theorem 7. If $f \in \mathcal{A}$ belongs to \mathcal{R}_{S_G} , then

$$|a_3a_5 - a_4^2| \leq \frac{146\,831}{3087\,360}. \quad (3.26)$$

Proof. From (3.13)–(3.15), we have

$$\begin{aligned} |a_3a_5 - a_4^2| &= \left| -\frac{29}{2949\,120}c_1^6 - \frac{1}{30\,720}c_1^4c_2 + \frac{3}{10\,240}c_1^3c_3 - \frac{1}{480}c_4c_1^2 \right. \\ &\quad \left. + \frac{7}{1920}c_1c_2c_3 - \frac{1}{480}c_2^3 + \frac{1}{240}c_4c_2 - \frac{1}{256}c_3^2 \right| \\ &\leq \frac{29}{2949\,120}|c_1|^6 + \frac{3}{10240}|c_1|^3 \left| c_3 - \frac{1}{9}c_1c_2 \right| + \frac{1}{240}|c_4| \left| c_2 - \frac{1}{2}c_1^2 \right| \\ &\quad + \frac{1}{256}|c_3| \left| c_3 - \frac{14}{15}c_1c_2 \right| + \frac{1}{480}|c_2|^3. \end{aligned}$$

Now using (2.3)–(2.5), we get the required result. \square

We will now determine the bound of the third Hankel determinant $H_{3,1}(f)$ for $f \in \mathcal{R}_{S_G}$.

Theorem 8. If $f \in \mathcal{A}$ belongs to \mathcal{R}_{S_G} , then

$$|H_{3,1}(f)| \leq \frac{319}{8640}. \quad (3.27)$$

Proof. We have the third Hankel determinant form as follows:

$$H_{3,1}(f) = a_3(a_2a_4 - a_3^2) - a_4(a_4 - a_2a_3) + a_5(a_3 - a_2^2) \quad (3.28)$$

where $a_1 = 1$. This yields

$$|H_{3,1}(f)| \leq |a_3||a_2a_4 - a_3^2| + |a_4||a_4 - a_2a_3| + |a_5||a_3 - a_2^2|. \quad (3.29)$$

By using (3.2)–(3.4) and (3.21)–(3.23), we obtain the desired result. \square

4. Bounds of $|H_{4,1}(f)|$ for the class \mathcal{R}_{S_G}

From (1.4), we can write $H_{4,1}(f)$ as

$$H_{4,1}(f) = a_7H_{3,1}(f) - a_6\delta_1 + a_5\delta_2 - a_4\delta_3, \quad (4.1)$$

where

$$\delta_1 = a_3(a_2a_5 - a_3a_4) - a_4(a_5 - a_2a_4) + a_6(a_3 - a_2^2), \quad (4.2)$$

$$\delta_2 = a_3(a_3a_5 - a_4^2) - a_5(a_5 - a_2a_4) + a_6(a_4 - a_2a_3), \quad (4.3)$$

$$\delta_3 = a_4(a_3a_5 - a_4^2) - a_5(a_2a_5 - a_3a_4) + a_6(a_2a_4 - a_3^2). \quad (4.4)$$

In recent years, researchers start to find the fourth-order Hankel determinant for different subclasses of analytic functions. The trend of finding fourth-order Hankel determinant in geometric function theory started in 2018, when Arif et al. [1] studied and successfully obtained the upper bound for the class of bounded turning functions. Recently Khan et al. [16] obtained the third and fourth-order Hankel determinant for the class of bounded turning functions associated with sine function. Also, Zhang and Tang [41] studied the fourth-order Hankel determinant for class of starlike functions connected with sine function. Inspired from the above works, we discuss here the fourth-order Hankel determinant for the class \mathcal{R}_{S_G} .

Theorem 9. *If $f \in \mathcal{A}$ belongs to \mathcal{R}_{S_G} , then*

$$|H_{4,1}(f)| \leq \frac{2334\ 260\ 186\ 533}{6535\ 323\ 648\ 000} \simeq 0.3571.$$

Proof. From (4.1), we have

$$H_{4,1}(f) = a_7H_{3,1}(f) - a_6\delta_1 + a_5\delta_2 - a_4\delta_3,$$

where δ_1 , δ_2 and δ_3 are defined as in (4.2)–(4.4). Now by using the triangle inequality, we have

$$|H_{4,1}(f)| \leq |a_7||H_{3,1}(f)| + |a_6||\delta_1| + |a_5||\delta_2| + |a_4||\delta_3|, \quad (4.5)$$

Since

$$\begin{aligned} |\delta_1| &= \left| a_3 (a_2 a_5 - a_3 a_4) - a_4 (a_5 - a_2 a_4) + a_6 (a_3 - a_2^2) \right| \\ &\leq |a_3| |a_2 a_5 - a_3 a_4| + |a_4| |a_5 - a_2 a_4| + |a_6| |a_3 - a_2^2|, \end{aligned}$$

by using (3.2), (3.3), (3.5), (3.21), (3.24) and (3.25), we get

$$|\delta_1| \leq \frac{3983}{17\,280}. \quad (4.6)$$

Since

$$\begin{aligned} |\delta_2| &= \left| a_3 (a_3 a_5 - a_4^2) - a_5 (a_5 - a_2 a_4) + a_6 (a_4 - a_2 a_3) \right| \\ &\leq |a_3| |a_3 a_5 - a_4^2| + |a_5| |a_5 - a_2 a_4| + |a_6| |a_4 - a_2 a_3|, \end{aligned}$$

by using (3.2), (3.4), (3.5), (3.22), (3.25) and (3.26), we get

$$|\delta_2| \leq \frac{15\,931\,363}{92\,620\,800}. \quad (4.7)$$

Also, since

$$\begin{aligned} |\delta_3| &= \left| a_4 (a_3 a_5 - a_4^2) - a_5 (a_2 a_5 - a_3 a_4) + a_6 (a_2 a_4 - a_3^2) \right| \\ &\leq |a_4| |a_3 a_5 - a_4^2| + |a_5| |a_2 a_5 - a_3 a_4| + |a_6| |a_2 a_4 - a_3^2|, \end{aligned}$$

by using (3.3)–(3.5), (3.23), (3.24) and (3.26), we get

$$|\delta_3| \leq \frac{53\,037\,859}{1111\,449\,600}. \quad (4.8)$$

Now by using the values of (4.6)–(4.8) along with (3.3)–(3.6) and (3.27) to (4.5), we get the desired estimate. \square

5. Bounds of $|H_{4,1}(f)|$ for the 2-fold and 3-fold symmetric functions

A function f is said to be m -fold symmetric if the following condition holds true for $\varepsilon = \exp\left(\frac{2\pi i}{m}\right)$,

$$f(\varepsilon z) = \varepsilon f(z) \quad (z \in \mathbb{D}).$$

The set of all m -fold symmetric functions belonging to the familiar class \mathcal{S} of univalent functions is denoted by $\mathcal{S}^{(m)}$, represented by the following series expansion

$$f(z) = z + \sum_{n=1}^{\infty} a_{mn+1} z^{mn+1} \quad (z \in \mathbb{D}). \quad (5.1)$$

An analytic function f of the form (5.1) belongs to the class $\mathcal{R}_{\mathcal{S}_G}^{(m)}$ if and only if

$$f'(z) = \frac{2}{1 + e^{-\left(\frac{p(z)-1}{p(z)+1}\right)}} \quad (z \in \mathbb{D}), \quad (5.2)$$

where $p(z)$ belong to the class $\mathcal{P}^{(m)}$ which is defined as follows:

$$\mathcal{P}^{(m)} = \left\{ p \in \mathcal{P} : p(z) = 1 + \sum_{n=1}^{\infty} c_{mn} z^{mn} \right\}. \quad (5.3)$$

If a function f belongs to $\mathcal{S}^{(2)}$, then its series representation is

$$f(z) = z + a_3 z^3 + a_5 z^5 + \dots,$$

and

$$H_{4,1}(f) = a_3 a_5 a_7 - a_3^3 a_7 + a_3^2 a_5^2 - a_5^3. \quad (5.4)$$

Further, if a function f belongs to $\mathcal{S}^{(3)}$, then its series representation is

$$f(z) = z + a_4 z^4 + a_7 z^7 + \dots,$$

and

$$H_{4,1}(f) = a_4^2 (a_4^2 - a_7). \quad (5.5)$$

Theorem 10. If $f \in \mathcal{R}_{S_G}^{(2)}$, then

$$|H_{4,1}(f)| \leq \frac{299}{108000}.$$

Proof. Let $f \in \mathcal{R}_{S_G}^{(2)}$. Then by the series (5.1)–(5.3) for $m = 2$, we have

$$\begin{aligned} f'(z) &= 1 + 3a_3 z^2 + 5a_5 z^4 + 7a_7 z^6 + \dots, \\ \frac{2}{1 + e^{-\left(\frac{c_2 z^2 + c_4 z^4 + \dots}{2 + c_2 z^2 + c_4 z^4 + \dots}\right)}} &= 1 + \frac{1}{4} c_2 z^2 + \left(\frac{1}{4} c_4 - \frac{1}{8} c_2^2\right) z^4 + \left(\frac{11}{192} c_2^3 - \frac{1}{4} c_4 c_2 + \frac{1}{4} c_6\right) z^6 + \dots. \end{aligned}$$

After comparing, we get

$$\begin{aligned} a_3 &= \frac{1}{12} c_2, \\ a_5 &= \frac{1}{20} \left(c_4 - \frac{1}{2} c_2^2 \right), \\ a_7 &= \frac{11}{1344} c_2^3 - \frac{1}{28} c_2 c_4 + \frac{1}{28} c_6. \end{aligned}$$

Then by substituting the above values to (5.4), we get

$$\begin{aligned} H_{4,1}(f) &= -\frac{529}{290\,304\,000} c_2^6 + \frac{437}{192\,000} c_2^4 c_4 - \frac{23}{241\,920} c_6 c_2^3 \\ &\quad + \frac{113}{2016\,000} c_2^2 c_4^2 + \frac{1}{6720} c_6 c_2 c_4 - \frac{1}{8000} c_4^3 \end{aligned}$$

and after rearranging, we get

$$H_{4,1}(f) = \frac{437}{24\,192\,000} c_2^4 \left(c_4 - \frac{23}{228} c_2^2 \right) + \frac{1}{6720} c_2 c_6 \left(c_4 - \frac{23}{36} c_2^2 \right) - \frac{1}{8000} c_4^3 \left(c_4 - \frac{113}{252} c_2^2 \right).$$

Now by using the triangle inequality along with (2.3) and (2.4), we get the required result. \square

Theorem 11. If $f \in \mathcal{R}_{S_G}^{(3)}$, then

$$|H_{4,1}(f)| \leq \frac{1}{896}.$$

Proof. Let $f \in \mathcal{R}_{S_G}^{(3)}$. Then by (5.1)–(5.3) for $m = 3$, we have

$$f'(z) = 1 + 4a_4z^3 + 7a_7z^6 + \dots \quad (5.6)$$

$$\frac{2}{1 + e^{-\left(\frac{c_3z^3 + c_6z^6 + \dots}{2 + c_3z^3 + c_6z^6 + \dots}\right)}} = 1 + \frac{1}{4}c_3z^3 + \left(\frac{1}{4}c_6 - \frac{1}{8}c_3^2\right)z^6 + \dots \quad (5.7)$$

After comparing (5.6) and (5.7), we get

$$\begin{aligned} a_4 &= \frac{1}{16}c_3, \\ a_7 &= \frac{1}{28}\left(c_6 - \frac{1}{2}c_3^2\right). \end{aligned}$$

Then by substituting the above values to (5.5), we get

$$H_{4,1}(f) = -\frac{1}{7168}c_3^2\left(c_6 - \frac{39}{64}c_3^2\right).$$

Now by using the (2.3) and (2.4), we get the required result. \square

6. Zalcman functional

One of the main conjectures in Geometric function theory, suggested by Lawrence Zalcman in 1960, is that the coefficients of class \mathcal{S} satisfy the inequality,

$$|a_n^2 - a_{2n-1}| \leq (n-1)^2. \quad (6.1)$$

Only the well-known Koebe function $k(z) = \frac{z}{(1-z)^2}$ and its rotations have equality in the above form. For the popular Fekete-Szegő inequality, when $n = 2$, the equality holds. Many researchers have researched Zalcman functional in the literature [5, 8, 19].

Theorem 12. Let $f \in \mathcal{A}$ belong to \mathcal{R}_{S_G} . Then

$$|a_3^2 - a_5| \leq \frac{1}{10}. \quad (6.2)$$

The result is sharp for the function f_4 defined by (3.7).

Proof. We use the Eqs (3.13) and (3.15) to get the Zalcman functional, and then we get

$$|a_3^2 - a_5| = \frac{1}{20} \left| \frac{37}{288}c_1^4 - \frac{119}{144}c_1^2c_2 + c_3c_1 + \frac{23}{36}c_2^2 - c_4 \right|.$$

Using Lemma 4, we can get the necessary result for the last expression. \square

7. Krushkal inequality for the class \mathcal{R}_{S_G}

In this section we will give a direct proof of the inequality

$$|a_n^p - a_2^{p(n-1)}| \leq 2^{p(n-1)} - n^p$$

over the class \mathcal{R}_{S_G} for the choice of $n = 4$, $p = 1$, and for $n = 5$, $p = 1$. Krushkal introduced and proved this inequality for the whole class of univalent functions in [17].

Theorem 13. *Let $f \in \mathcal{A}$ belong to \mathcal{R}_{S_G} . Then*

$$|a_4 - a_2^3| \leq \frac{1}{8}.$$

The result is sharp for the function f_3 defined by (3.7).

Proof. From Eqs (3.12) and (3.14), we get

$$|a_4 - a_2^3| = \left| \frac{19}{1536}c_1^3 - \frac{1}{16}c_2c_1 + \frac{1}{16}c_3 \right|.$$

By applying (2.6) to the above, we get the required result. \square

Theorem 14. *Let $f \in \mathcal{A}$ belong to \mathcal{R}_{S_G} . Then*

$$|a_5 - a_2^4| \leq \frac{1}{10}.$$

The result is sharp for the function f_4 defined by (3.7).

Proof. From Eqs (3.12) and (3.14), we get

$$|a_5 - a_2^4| = \frac{1}{20} \left| \frac{101}{1024}c_1^4 - \frac{11}{16}c_1^2c_2 + c_3c_1 + \frac{1}{2}c_2^2 - c_4 \right|.$$

By using Lemma 4, we can get the necessary result for the last expression. \square

8. Conclusions

In the present study, we have defined the class of bounded turning functions associated with modified sigmoid function. Also we have determined the sharp results for some coefficient functionals which play a very important role in the study of the geometric function theory. Furthermore, we have evaluated bounds of the third and fourth-order Hankel determinants for the 2-fold and 3-fold symmetric functions.

Recently, the usages of the quantum (or q -) calculus happens to provide another popular direction for researches in geometric function theory of complex analysis. This is evidenced by the recently-published survey-cum-expository review article by Srivastava [31]. Therefore the quantum (or q -) extensions of the results, which we have presented in this paper, are worthy of investigation.

Acknowledgements

The authors would like to express their gratitude to the editor and the reviewers for their valuable comments. The second author was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (No. 2019R1I1A3A01050861).

Conflict of interest

The authors declare that they have no competing interests.

References

1. M. Arif, L. Rani, M. Raza, P. Zaprawa, Fourth Hankel determinant for the family of functions with bounded turning, *Bull. Korean Math. Soc.*, **55** (2018), 1703–1711. doi: 10.1007/s40278-018-49036-z.
2. M. Arif, M. Raza, H. Tang, S. Hussain, H. Khan, Hankel determinant of order three for familiar subsets of analytic functions related with sine function, *Open Math.*, **17** (2019), 1615–1630. doi: 10.1515/math-2019-0132.
3. K. O. Babalola, On $H_3(1)$ Hankel determinant for some classes of univalent functions, *Inequal. Theory. Appl.*, **6** (2017), 1–7.
4. O. M. Barukab, M. Arif, M. Abbas, S. A. Khan, Sharp bounds of the coefficient results for the family of bounded turning functions associated with a Petal-shaped domain, *J. Funct. Spaces*, **2021** (2021), 5535629. doi: 10.1155/2021/5535629.
5. J. E. Brown, A. Tsao, On the Zalcman conjecture for starlikeness and typically real functions, *Math. Z.*, **191** (1986), 467–474. doi: 10.1007/BF01162720.
6. N. E. Cho, V. Kumar, S. Kumar, V. Ravichandran, Radius problems for starlike functions associated with the sine function, *Bull. Iran. Math. Soc.*, **45** (2019), 213–232. doi: 10.1007/s41980-018-0127-5.
7. N. E. Cho, S. Kumar, V. Kumar, V. Ravichandran, H. M. Srivastava, Starlike functions related to the Bell numbers, *Symmetry*, **11** (2019), 219. doi: 10.3390/sym11020219.
8. D. Bansal, J. Sokól, Zalcman conjecture for some subclass of analytic functions, *J. Fractional Calculus Appl.*, **8** (2017), 1–5.
9. J. Dziok, R. K. Raina, R. K. J. Sokól, On a class of starlike functions related to a shell-like curve connected with Fibonacci numbers, *Math. Comput. Model.*, **57** (2013), 1203–1211. doi: 10.1016/j.mcm.2012.10.023.
10. P. Goel, S. S. Kumar, Certain class of starlike functions associated with modified sigmoid function, *Bull. Malays. Math. Sci. Soc.*, **43** (2020), 957–991. doi: 10.1007/s40840-019-00784-y.
11. W. K. Hayman, On the second Hankel determinant of mean univalent functions, *Proc. London Math. Soc.*, **3** (1968), 77–94. doi: 10.1112/plms/s3-18.1.77.

12. S. Kanas, D. Răducanu, Some classes of analytic functions related to conic domains, *Math. Slovaca*, **64** (2014), 1183–1196. doi: 10.2478/s12175-014-0268-9.
13. F. R. Keogh, E. P. Merkes, A coefficient inequality for certain classes of analytic functions, *Proc. Amer. Math. Soc.*, **20** (1969), 8–12. doi : 10.1017/s0084255900011207.
14. W. Keopf, On the Fekete-Szegő problem for close-to-convex functions, *Proc. Amer. Math. Soc.*, **101** (1987), 89–95.
15. M. G. Khan, B. Ahmad, G. M. Moorthy, R. Chinram, W. K. Mashwani, Applications of modified Sigmoid functions to a class of starlike functions, *J. Funct. Spaces*, **2020** (2020), 8844814.
16. M. G. Khan, B. Ahmad, J. Sokól, Coefficient problems in a class of functions with bounded turning associated with sine function, *Eur. J. Pure. Appl. Math.*, **14** (2021), 53–64. doi: 10.51202/0323-3243-2021-3-053.
17. S. K. Krushkal, A short geometric proof of the Zalcman and Bieberbach conjectures, *arXiv*. Available from: <https://arxiv.org/abs/1408.1948>.
18. R. J. Libera, E. J. Zlotkiewicz, Early coefficients of the inverse of a regular convex function, *Proc. Amer. Math. Soc.*, **85** (1982), 225–230. doi: 10.1090/S0002-9939-1982-0652447-5.
19. W. C. Ma, The Zalcman conjecture for close-to-convex functions, *Proc. Amer. Math. Soc.*, **104** (1988), 741–744. doi: 10.1090/S0002-9939-1988-0964850-X
20. W. C. Ma, D. Minda, A unified treatment of some special classes of univalent functions, *Proceedings of the conference on complex analysis*, Tianjin, 1992.
21. R. Mendiratta, S. Nagpal, V. Ravichandran, On a subclass of strongly starlike functions associated exponential function, *Bull. Malays. Math. Sci. Soc.*, **38** (2015), 365–386. doi: 10.1093/oxartj/kcv022.
22. J. W. Noonan, D. K. Thomas, On the Second Hankel determinant of a really mean p -valent functions, *Trans. Amer. Math. Soc.*, **22** (1976), 337–346. doi: 10.1090/S0002-9947-1976-0422607-9.
23. H. Orhan, N. Magesh, J. Yamini, Bounds for the second Hankel determinant of certain bi-univalent functions, *Turkish J. Math.*, **40** (2016), 679–687. doi: 10.3906/mat-1505-3.
24. C. Pommerenke, On the Hankel determinants of univalent functions, *Mathematika*, **14** (1967), 108–112. doi: 10.1112/S002557930000807X.
25. C. Pommerenke, *Univalent functions*, Göttingen: Vanderhoeck & Ruprecht, 1975.
26. R. K. Raina, J. Sokól, On coefficient estimates for a certain class of starlike functions, *Hacettepe. J. Math. Statist.*, **44** (2015), 1427–1433.
27. V. Ravichandran, S. Verma, Bound for the fifth coefficient of certain starlike functions, *Comptes Rendus Math.*, **353** (2015), 505–510. doi: 10.1016/j.crma.2015.03.003.
28. L. Shi, M. G. Khan, B. Ahmad, W. K. Mashwani, P. Agarwal, S. Momani, Certain coefficient estimate problems for three-leaf-type starlike functions, *Fractal Fract.*, **5** (2021), 137. doi: 10.3390/fractalfract5040137.
29. K. Sharma, N. K. Jain, V. Ravichandran, Starlike functions associated with cardioid, *Afr. Mat.*, **27** (2016), 923–939. doi: 10.1007/s13370-015-0387-7.

30. J. Sokół, S. Kanas, Radius of convexity of some subclasses of strongly starlike functions, *Zesz. Nauk. Politech. Rzeszowskiej Mat.*, **19** (1996), 101–105.
31. H. M. Srivastava, Operators of basic (or q -) calculus and fractional q -calculus and their applications in geometric function theory of complex analysis, *Iran. J. Sci. Technol. Trans. A: Sci.*, **44** (2020), 327–344. doi: 10.1007/s40995-019-00815-0.
32. H. M. Srivastava, Q. Z. Ahmad, M. Darus, N. Khan, B. Khan, N. Zaman, et al., Upper bound of the third Hankel determinant for a subclass of close-to-convex functions associated with the lemniscate of Bernoulli, *Mathematics*, **7** (2019), 848. doi: 10.3390/math7090848.
33. H. M. Srivastava, Ş. Altinkaya, S. Yalçın, Hankel determinant for a subclass of bi-univalent functions defined by using a symmetric q -derivative operator, *Filomat*, **32** (2018), 503–516. doi: 10.2298/FIL1802503S.
34. L. Shi, M. G. Khan, B. Ahmad, Some geometric properties of a family of analytic functions involving a generalized q -operator, *Symmetry*, **12** (2020), 291. doi: 10.3390/sym12020291.
35. H. M. Srivastava, Some parametric and argument variations of the operators of fractional calculus and related special functions and integral transformations, *J. Nonlinear Convex Anal.*, **22** (2021), 1501–1520.
36. H. M. Srivastava, B. Khan, N. Khan, M. Tahir, S. Ahmad, N. Khan, Upper bound of the third Hankel determinant for a subclass of q -starlike functions associated with the q -exponential function, *Bull. Sci. Math.*, **167** (2021), 102942. doi: 10.1016/j.bulsci.2020.102942.
37. H. M. Srivastava, G. Kaur, G. Singh, Estimates of the fourth Hankel determinant for a class of analytic functions with bounded turnings involving cardioid domains, *J. Nonlinear Convex Anal.*, **22** (2021), 511–526.
38. M. Shafiq, H. M. Srivastava, N. Khan, Q. Z. Ahmad, M. Darus, S. Kiran, An upper bound of the third Hankel determinant for a subclass of q -starlike functions associated with k -Fibonacci numbers, *Symmetry*, **12** (2020), 1043. doi: 10.3390/sym12061043.
39. L. A. Wani, A. Swaminathan, Starlike and convex functions associated with a Nephroid domain, *Bull. Malays. Math. Sci. Soc.*, **44** (2021), 79–104. doi: 10.1007/s40840-020-00935-6.
40. D. M. Wilkes, S. D. Morgera, F. Noor, M. H. Hayes, A Hermitian Toeplitz matrix is unitarily similar to a real Toeplitz-plus-Hankel matrix, *IEEE Trans. Signal Process.*, **39** (1991), 2146–2148. doi: 10.1109/78.134459
41. H. Y. Zhang, H. Tang, A study of fourth-order Hankel determinants for starlike functions connected with the sine function, *J. Funct. Spaces*, **2021** (2021), 9991460. doi: 10.1155/2021/9991460.



AIMS Press

©2022 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)