

Sustainable Agriculture Reviews 22

Eric Lichtfouse *Editor*

Sustainable Agriculture Reviews

 Springer

Sustainable Agriculture Reviews

Volume 22

Series editor

Eric Lichtfouse

Other Publications by Dr. Eric Lichtfouse

Books

Scientific Writing for Impact Factor Journals
Nova Publishers 2013

Sustainable Agriculture
Springer 2009

Sustainable Agriculture Volume 2
Springer 2011

Environmental Chemistry. Green Chemistry and Pollutants in Ecosystems
Springer 2005

Rédiger pour être publié ! Conseils pratiques pour les scientifiques
Springer 2012, 2^e édition.

Journals and Series

Agronomy for Sustainable Development
www.springer.com/journal/13593

Sustainable Agriculture Reviews
www.springer.com/series/8380

Environmental Chemistry Letters
www.springer.com/journal/10311

Environmental Chemistry for a Sustainable World
www.springer.com/journal/11480

Blog

Agronomy blog
<http://www1.montpellier.inra.fr/agronomy-blog>

Magazine

Publier La Science
https://listes.inra.fr/sympa/d_read/veillecaps/

More information about this series at <http://www.springer.com/series/8380>

Eric Lichtfouse
Editor

Sustainable Agriculture Reviews



Editor

Eric Lichtfouse

CEREGE

INRA

Aix en Provence, France

ISSN 2210-4410

Sustainable Agriculture Reviews

ISBN 978-3-319-48005-3

DOI 10.1007/978-3-319-48006-0

ISSN 2210-4429 (electronic)

ISBN 978-3-319-48006-0 (eBook)

Library of Congress Control Number: 2016963152

© Springer International Publishing Switzerland 2017

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

This Springer imprint is published by Springer Nature

The registered company is Springer International Publishing AG

The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

Contents

1	Animal Waste: Opportunities and Challenges	1
	Francesca Girotto and Raffaello Cossu	
2	Soil Microorganisms Can Reduce P Loss from Cropping Systems	15
	R. Michael Lehman and Wendy I. Taheri	
3	Greenhouse Technology for Agriculture Under Arid Conditions	37
	Ellen Letsogile Thipe, Tilahun Workneh, Alferd Odindo, and Mark Laing	
4	Biochar for Agriculture in Pakistan	57
	Fahd Rasul, Ashfaq Ahmad, Muhammad Arif, Ishaq Ahmad Mian, Kawsar Ali, Muhammad Farooq Qayyum, Qaiser Hussain, Muhammad Aon, Shahzad Latif, Ruben Sakrabani, Muhammad Saghir, Genxing Pan, and Simon Shackley	
5	Effect of UV-B Radiation on Leguminous Plants.....	115
	Krishna Kumar Choudhary and Shashi Bhushan Agrawal	
6	Control of Fungal Diseases in Agricultural Crops by Chitinase and Glucanase Transgenes.....	163
	Jagdeep S. Sandhu, Manpreet K. Sidhu, and Inderjit S. Yadav	
7	Bioengineering Hairy Roots: Phytoremediation, Secondary Metabolism, Molecular Pharming, Plant-Plant Interactions and Biofuels	213
	Peyman Habibi, Maria Fatima Grossi De Sa, Abdullah Makhzoum, Sonia Malik, André Luís Lopes da Silva, Kathleen Hefferon, and Carlos Ricardo Soccol	

8	Soil Management to Optimize Water in Rice-Wheat Cropping	253
	Jagdish Singh, Sandeep Singh Sandhu, Dhanwinder Singh, and Manjit Singh Hadda	
9	Microbial Inoculants for Soil Quality and Plant Health	281
	Elizabeth T. Alori, Michael O. Dare, and Olubukola O. Babalola	
10	Conservation Agriculture in Tanzania	309
	Msafiri Yusuph Mkonda and Xinhua He	
	Index.....	325

Chapter 9

Microbial Inoculants for Soil Quality and Plant Health

Elizabeth T. Alori, Michael O. Dare, and Olubukola O. Babalola

Abstract Agriculture is the major economic activity of most developing countries engaging more than 50 % of the population. Low world crop productivity due to low soil moisture, low nutrient capital, erosion risk, low pH, high phosphorus fixation, low levels of soil organic matter, aluminum toxicity pest and diseases, weeds and loss of soil biodiversity has induced the green revolution agriculture which involves high yielding varieties and agrochemicals. The continuous use of fertilizers, pesticides and herbicides has led to low agricultural productivity, low soil fertility, unfavourable economic returns, food poisoning, soil damage loss of biodiversity and serious environmental hazards. Microbial inoculants possess the capacity to enhance nutrient availability, uptake, and support the health of soil and plants to promote sustainable yield and has therefore gained attention of many agriculturist and researchers.

We review the ability of soil through the use of microbial inoculants to supply nitrogen, phosphorus and potassium to crop plants and enhance structural stability. Microbial inoculants such as rhizobium, plant growth promoting rhizobacteria and arbuscular mycorrhizal fungi can be used as biofertilizer to improve soil nitrogen, phosphorus and potassium availability and uptake. Both bacteria and fungi inoculants show potential for use in soil aggregate formation and stabilization and hence,

E.T. Alori

Department of Crop and Soil Sciences, Landmark University, Omu-Aran, Nigeria

Food Security and Safety Niche Area, Faculty of Agriculture, Science and Technology,
2735 Mmabatho, Mafikeng, South Africa

e-mail: alori.elizabeth@lmu.edu.ng

M.O. Dare

Food Security and Safety Niche Area, Faculty of Agriculture, Science and Technology,
2735 Mmabatho, Mafikeng, South Africa

O.O. Babalola (✉)

Food Security and Safety Niche Area, Faculty of Agriculture, Science and Technology,
2735 Mmabatho, Mafikeng, South Africa

Food Security and Safety Niche Area, Faculty of Agriculture, Science and Technology,
North-West University, 2735 Mmabatho, Mafikeng, South Africa

e-mail: olubukola.babalola@nwu.ac.za

soil structure enhancement. The ability of microbial inoculants to ameliorate plant stress as a result of drought, soil contamination and salinity are also highlighted. The most commonly used microorganisms as biofertilizers, biocontrol and bioremediators include *Bacillus* spp, *Pseudomonas* spp, *Streptomyces* spp *Trichoderma* spp and *Mycorrhizas*. Microbial inoculants function through various mechanisms such as production of plant hormones, expansion and elongation of the root system, eliciting induced systemic resistance or systemic acquired resistance, production of lytic enzyme and antibiotic 4-hydroxyphenylactic acid, and production of 1-amino cyclopropane-1-carboxylate-deaminase (ACC-deaminase) in plants rhizosphere. These strategies are safe and sustainable in the long run. The use of appropriate carrier material determines the success of microbial inoculation techniques. Microbial inoculants could either be applied directly to the soil or as seed dressing. The fate of microbial inoculants under field application depends largely on both biotic and abiotic factors. The application of some microbial inoculants could cause a change (which could be a decrease or an increase) in the equilibrium of soil microbial communities while some produce no effect at all.

Keywords Agricultural sustainability • Biocontrol • Biofertilizer • Bioremediation • Biotechnology • Food security • Microbial inoculants • Plant growth • Plant growth promoting microorganisms (PGPM) • Soil fertility and health

9.1 Introduction

The increasing demand for food production with shrinking land resources is a major challenge to agricultural sustainability. Sustainable food production requires efficient use of determinate resources (Owen et al. 2015). Attempt to mitigate the problem include the use of high yielding varieties, chemical fertilizers and pesticides to supplement plant nutrition and control plant pathogens for increased agricultural productivity. However the increasing impacts of these agricultural practices on the environment have gradually affected the quality of soil hence, there is a need to optimize soil productivity in such a way that soil capacity to function as a healthy medium is preserved (Trivedi et al. 2012). The use of eco-friendly resources or input has been a major focus of attention in the past three decades. Although reports on the benefits of using microbial inoculants for plant growth promotion and health in agricultural soil have been inconsistent, there is a promising trend for microbial inoculants to meet the sustainable agricultural production needs. Suggestions to replace or supplement the heavy application of chemical fertilizers with inoculants have been reported (Carvajal-Muñoz and Carmona-Garcia 2012). Microbial inoculants application has been in existence for more than 100 years but gained a lot of prominence in the last three decades with several commercial inoculants products in the market (Babalola and Glick 2012).




Fig. 9.1 Maize plant (a). Showing *Fusarium graminearum* infection (b). Inoculated with *Pseudomonas* sp for biocontrol against *Fusarium graminearum*

Microbial inoculants participate in many ecosystem biological and chemical processes such as biological control of pathogens (Fig. 9.1) and nutrient cycling, thereby improving nutrient availability. Microbial inoculants application increase biodiversity, creating suitable condition for development of beneficial microorganism. They also improve physical properties of soil such as; improve structure and aggregation of soil particles; reduce soil compaction, increase spore spaces and water infiltration. The antioxidant properties of microbial inoculants promote decomposition of organic matter and increase humus content in soil matrix, and are therefore being considered as an alternative way of reducing the use of chemicals in agriculture (Carvajal-Muñoz and Carmona-Garcia 2012). Microbial inoculants techniques ensure biodegradation of complex substances and develop bioremediation processes in soil contaminated with toxics, xenobiotic and recalcitrant substances.

The strategies involved in plant growth promotion by microbial inoculants could be a direct or indirect mechanism. Directly, inoculation of crop plant with microbial inoculants could result in the expansion and elongation of the root system, leading to improved uptake of water and nutrients (Halpern et al. 2015). Production of growth hormones by microbial inoculants impact root morphogenesis such that plant root hairs and lateral roots are over produced resulting in greater uptake of plant nutrients and hence improvement of plant growth (Kumar et al. 2007). Fixations of atmospheric nitrogen, solubilization of minerals such as phosphorus (P) (Babalola 2010), are also some of the direct mechanisms of influence of microbial inoculants. In indirect growth promotion, Microbial inoculants affect the status of plants by eliciting induced systemic resistance (ISR) or systemic acquired resistance (SAR), by improving disease resistance. These acts prevent soil-borne pathogens from inhibiting plant growth (Yang et al. 2009). Ability to trigger a salicylic acid (SA) -independent pathway controlling systemic resistance is a common trait

of ISR-inducing biocontrol bacteria. Structural deformities in pathogenic fungi under in vitro culture conditions by the production of diffusible and volatile antifungal compounds have been reported. The bacterial strain successfully restricted the growth of all the test fungi in dual cultures and induced morphological abnormalities such as mycelial and conidial deviations. Also of note is the production of siderophores that solubilize and sequester iron (Hmaeid et al. 2014).

With the increasing use of microbial inoculants for plant growth promotion, this review discusses some of the beneficiary roles of microbial inoculants in plant and soil. It describes changes in soil structure, nutrient solubility as a result of the application of microbial inoculants. We provide an overview of microbial inoculants use for agricultural sustainability, the significance of their application on soil nutrient improvement and soil structure enhancement. Their roles in amelioration of plants stress as a result of drought, soil contaminants, salinity and as biocontrol agents are well explained.

9.2 Microbial Inoculants

Microbial inoculation is one of the major agricultural practices that have been used to acquire desirable characteristics in the soil. Microbial inoculants are the formulations of beneficial living microorganisms that when added to soil, improve availability of nutrient to host plant directly or indirectly, thereby promoting plant growth (Gaiad 2011). Most of the microorganisms that are used in the production of microbial inoculants inhabit or are capable of inhabiting the soil and perform various roles and functions in the soil. Microbial inoculants are applied, singly or in combinations, to seeds, plants and soil to enhance their productivity. Different terminologies such as biostimulant (Halpern et al. 2015), bio-inoculants (Singh et al. 2013), and bio-fertilizers (Ansari et al. 2014) have been used to represent these groups of microorganisms. Microbial inoculants include three major groups: (1) plant growth promoting rhizobacteria (PGPR), (2) arbuscular mycorrhiza fungi and (3) the nitrogen-fixing *rhizobia*, which are usually not considered as PGPR (Yadav and Verma 2014). These groups are known to possess the capacity to enhance nutrient availability, uptake, and support the health of plants to promote plant growth. Microbial inoculants are not nutrients but microorganisms that are able to increase the availability of these nutrients through their biophysical and biochemical activities in soil. Business Communication Co. research report (2011) estimated a compound annual growth rate of about 6.9 % for global microbial inoculants with a market value of \$4.5 billion in 2010, \$4.9 billion in 2011 and projected to reach \$6.8 billion by 2016 (Chatzipavlidis et al. 2013).

Microbial inoculation provides an innovative and cost-effective alternative to overcome salinity stress in soils (Tank and Saraf 2010). The system is environmentally friendly, and poses no health risk to either plant, human or animal. Enhance soil nutrient availability to the plants hence their use as biofertilizers (Ahmad and Kibret 2014). Microbial inoculants provide resistance against pathogens. They can

Table 9.1 Examples of some microbial inoculants, the test crops and their beneficial properties

Microbial inoculants	Test Crop	Beneficial properties	References
<i>Chryseobacterium indologenes</i> , <i>Pseudomonas cepacia</i> , <i>P. fluorescens</i>	Wide barley	Salt stress	Hmaeid et al. (2014)
<i>Bacillus subtilis</i> , <i>P. corrugate</i>	Maize	Cool regions (22 °C)	Trivedi et al. (2012)
<i>Paenibacillus yonginensis</i> DCY84	<i>Arabidopsis thaliana</i>	Drought, salt stress	Sukweenadhia et al. (2015)
<i>Enterobacter sakazakii</i>	Cowpea	Parasitic weed	Babalola et al. (2007)
<i>Bacillus subtilis</i>	Cotton	Phytopathogen	Pereg and McMillan (2015)
<i>Scutellospora reticulate</i> , <i>Glomus pansihalos</i> (Mycorrhizal fungi)	Cowpea	Soil polluted with Al and Mn	Alori and Fawole (2012)
<i>Pseudomonas putida</i>	Wheat	Cool region	Trivedi and Pandey (2007)
<i>Trichoderma</i> sp., <i>Gliocladium</i> sp.	Flowers, ornamentals	Plant pathogens	Julia et al. (2013)
<i>Azotobacter</i> sp + <i>Pseudomonas</i> sp	Mustard	Cadmium	Panwar et al. (2011)
Arbuscular mycorrhizal fungi	Citrus	Drought stress	Wu et al. (2013)
<i>Burkholderia cepacia</i>	Yellow lupine	Toluene	Barac et al. (2009)
<i>P. fluorescens</i> strains, CHA0 and Pf1	Banana	Drought stress	Kavino et al. (2010)
<i>Gordonia</i> sp. S2Rp-17	Corn	Diesel (Soil contaminant)	Hong et al. (2011)
<i>Sinorhizobium meliloti</i>	Common reed	Phenanthrene	Golubev et al. (2009)
<i>Bacillus thuringiensis</i> , <i>Rhizophagus intraradices</i>	Trifolium repens	Drought stress	Ortiz et al. (2015)
<i>Burkholderia cepacia</i>	Popular	Toluene	Taghavi et al. (2005)
Arbuscular mycorrhizal	Olive	Salinity stress	Porras-Soriano et al. (2009)
<i>Azospirillum lipoferum</i>	Wheat	Crude oil	Muratova et al. (2005)

therefore be used in biological control against plant pathogens (Sukweenadhia et al. 2015) against weed pest (Biological herbicides) (Babalola et al. 2007) and insect pest (Saharan and Nehra 2011). Microbial inoculants can also be used in phytoremediation of polluted soils (Alori and Fawole 2012; Alori 2015). Waterlogged, compacted, desiccated wind and rain eroded soil are remediated through microbial inoculation. Fungal inoculants protect plants against transplant shock, promote environmental resistance to heat and drought (Sukweenadhia et al. 2015) and vastly improve the quality of the soil (Table 9.1).

9.2.1 Microbial Inoculants and Soil Fertility Improvement

9.2.1.1 Nitrogen (N)

N remains the most limiting element for plant growth. The major sources of N for agricultural soil are mineral fertilizers and biological N fixation carried out by microorganisms. Nitrogen-fixation is the first step for cycling N to the biosphere from the atmosphere, a key input of N for plant productivity (Bernhard 2010). Microbes especially bacteria are important in N cycling. Bacteria are known to exclusively fix atmospheric N either symbiotically or asymbiotically due to their possession of the key enzyme nitrogenase which specifically reduces atmospheric N to ammonia (Wagner 2011).

Symbiotic N fixation in soil is a process occurring in legume and non-legume plants. The bacteria *Rhizobium*, *Sinorhizobium*, *Allorhizobium*, *Bradyrhizobium*, *Mesorhizobium* and *Azorhizobium*, collectively referred to as *rhizobia* are responsible for the legume N fixation while *Frankia* and *Actinobacteria* are responsible for non-legume N fixation in soil (Wagner 2011). It is evident that inoculation of legumes with rhizobia has the ability to increase the soil N status. N fixed annually by legume-rhizobium association was reported to be about 40–48 million tonnes compared to 98 million t year⁻¹ of N fertilizer (Jenkinson 2001). This ability to fix high amounts of N into the soil is a great potential of rhizobial inoculant to reduce the cost of industrial N fertilizers, thereby reducing the cost of inputs for farmers. Nitrogen fixation of an effectively nodulated legume is a vital and indispensable aspect of sustainable agriculture. The use of rhizobium inoculant to achieve efficient N fixation in soil requires the compatibility of the legume and rhizobium inoculant and their adaptability to the environment (Wagner 2011). The few inconsistencies about the ability of the rhizobium inoculant to increase the soil N are probably due to compatibility and adaptability to the environment issue. Rhizobium must be able to establish, compete and persist with other microflora to form effective nodules in the introduced environment (Gaiad 2011). *Rhizobial* inoculation has played a vital role in legume production in the US and Australia. The legume-cereal cropping system popularized by the International Institute of Tropical Agriculture in sub-Saharan African is also an indication of the growing popularity of the use of *rhizobial* inoculant. Most crop rotation systems use legume in crop sequence because of the understanding that legumes are able to fix N and increase the N status of the soil for the next crop. This has been utilized significantly in soil fertility management for crops to reduce the application of chemical fertilizers. Therefore the proper and efficient use of *rhizobial* inoculants will ultimately benefit sustainable agricultural production.

A group of bacteria that are free living in the soil commonly referred to as plant growth promoting rhizobacteria/plant growth promoting bacteria/plant growth promoting microorganism have the ability to fix N into soil (Calvo et al. 2014), when occupying the rhizosphere of crops both legumes and non-legume. These groups of bacteria include the genera of *Pseudomonads*, *Azoarcus*, *Beijerinckia*, *Cyanobacteria*

(*Nostoc* and *Anabaena*), *Klebsiella*, *Pantoea*, *Azotobacter*, *Azospirillum*, *Bacillus*, *Burkholderia*, *Herbaspirillum*, and *Gluconacebacter diazotrophicus* (Egamberdiyeva 2007). Many of these organisms are used as microbial inoculants for crop growth improvement, singly or in combination with other organisms. Unlike rhizobial inoculants, the use of many of these bacteria as a single inoculant to increase soil N fertility for crop use has not been very effective. Although they may not be as effective as *rhizobium*, there is potential for improvement in their ability to help in sustainable agricultural production. Although the ability of many plant growth promoting rhizobacterial to increase N content of the soil are very inconsistent, some cases of appreciable soil N increase have been observed especially when the inoculants contain more than one of the organisms. The use of *Azospirillum* as helper bacteria used in combination with *rhizobium* increased the effect of *rhizobium* in soil fixation. *Azotobacter* and *rhizobium* contributed 78.8 kg N ha⁻¹ year⁻¹ total N to soil in soybean-wheat rotation (Rawat et al. 2013).

Plant nutrition for N has also been improved by the application of some fungal inoculants. Arbuscular mycorrhizal fungi that form associations with more than 80 % of plants including most crops have also been identified as a probable N mobilizer for plants (Hodge and Storer 2015; Veresoglou et al. 2012). However, the contribution of arbuscular mycorrhizal fungi to plant N uptake varies widely and the reasons for the variability are still unclear and may likely be resolved by the application of genomics and metabolomics technology (Hodge and Storer 2015). The role of arbuscular mycorrhizal fungi in plant P nutrition and soil structure improvement is well established, and the prospect of involving it in the N nutrition of crops will be a giant stride in tackling soil degradation problems through the use of microbial inoculants. Microbial inoculants such as rhizobium, plant growth promoting rhizobacteria and arbuscular mycorrhizal fungi have the potential to be used as biofertilizer to improve soil N availability and supply for sustainable agriculture

9.2.1.2 Phosphorus

Phosphorus is an essential macronutrient required by plants for their growth and development. It makes up about 0.1 % of the earth's crust (Sanderson 2014). However, most of the P in the earth's crust is in insoluble form and not readily available to plants. Low soil available P limits about 40 % of crop production in arable land worldwide (Bargaz et al. 2012). To compound the problem of P availability, added P fertilizers undergo fixation due to the complex exchanges within the soil limiting the availability of P to plants (Zhu et al. 2011).

The role of microbial inoculants in increasing the availability of soil P for plant growth can be viewed from two perspectives: firstly the solubilization of P from the mineral rock thereby increasing the available P in soil solution and secondly the mobilization of the available P to the plant roots for uptake. Phosphate solubilizing microorganisms and phosphate mobilizing microorganisms (Owen et al. 2015) include the genera of some bacteria and fungi that have been identified to solubilize and render insoluble soil P available to plants with their production of organic acids

and enzyme phosphatases and phytase (Calvo et al. 2014). These organisms include *Pseudomonas*, *Azospirillum*, *Azotobacter*, *Bacillus*, *Burkholderia*, *Enterobacter*, *Rhizobium*, *Erwinia*, *Streptomyces*, *Achromobacter* *Flavobacterium* and mycorrhiza (Ma et al. 2009). These microorganisms produce organic acids that chelate the cations bound to phosphate and convert them to soluble form (Calvo et al. 2014). They also produce the enzymes phytases and phosphatases that dephosphorylate phytates, the predominant organic P in the soil (60 % of organic P) to release P in a form available to plants (Singh and Satyanarayana 2011). The application of these organisms, either as bacterial or fungal inoculants, has advantage over the P fertilizers that readily form complexes in the soil when applied because the microbes can continuously supply available P to plants over a long range of time. Soil management practices that incorporate microbial inoculant application can really benefit from the sustained P supply to crops.

Fungal inoculants that have P solubilizing/mobilizing potential are well known. The most studied among the P mobilizing fungi is the arbuscular mycorrhizal fungi. Arbuscular mycorrhizal fungi are widespread in the plant kingdom and contribute significantly to plant P nutrition and growth in natural ecosystems (Smith et al. 2011). The mechanism of increased P uptake by Arbuscular mycorrhizal fungi has been attributed to the fungal extra radical hyphae growing beyond the phosphate depletion zone that develops around the root (Smith and Read 2008). Positive effects of arbuscular mycorrhizal fungi inoculation on the growth and P nutrition of crops have been reported (Cozzolino et al. 2013; Dare et al. 2010). Many mycorrhizal inoculants have been produced on a commercial scale, mostly in the US and Europe. *Rhizophagus* (formerly *Glomus*) *intraradices* and *Funneliformis* (formerly *Glomus*) *mosseae* (Kruger et al. 2012) are some of the common mycorrhizal inoculants which have been shown to increase P uptake in diverse crop plants (Dare et al. 2010; Cozzolino et al. 2013). Some other fungi such as *Aspergillus* and *Penicillium* species are able to solubilise inorganic phosphate and mineralise organic phosphate by secreting organic acids and producing phosphatase enzymes (Wang et al. 2015). The significant role of microbial inoculants in increasing the sustainable availability of P to plant is that of P solubilization and P mobilization.

9.2.1.3 Potassium (K)

Potassium is one of the most important macronutrient for plant growth and the third in fertilizer formulation after N and P. Potassium is one of the seven most common elements in the earth's crust and makes up 2.6 % of the earth's surface layer (Meena et al. 2014). Inadequate supplies of K to plants can lead to poor root growth, slow growth and lower yields in crops (White and Karley 2010).

K-solubilizing microorganisms present in soil and plant rhizosphere are evidently involved in the K cycles (Liu et al. 2012). Potassium-solubilizing microorganisms improve soil nutrients and structure and plant growth by releasing K from insoluble minerals into the soil (Meena et al. 2014). The microorganisms in the

soil or rhizosphere solubilize mineral K by synthesizing organic acids (Parmar and Sindhu 2013). A wide range of rhizospheric microorganisms that have been used as inoculants for increasing the K soil content or K plant nutrition include *Acidithiobacillus ferrooxidans*, *Arthrobacter* sp., *Bacillus edaphicus*, *Bacillus circulans*, *Bacillus mucilaginosus*, *Burkholderia* sp., and *Paenibacillus* sp. (Zarjani et al. 2013; Sangeeth et al. 2012). These organisms convert insoluble or mineral structural K compounds into soluble form and make them available for plants.

AM fungal inoculants have also been reported to increase K uptake. Arbuscular mycorrhizal releases proton H^+ or CO_2 and organic anions such as citrate malate and oxalate which increase the solubility of mineral K (Meena et al. 2014). However, the increased K by mycorrhizal has often been linked to increased P availability (Cardoso and Kuyper 2006). Plant growth promoting rhizobacteria and Arbuscular mycorrhizal fungi are responsible for K solubilization and mobilization for sustainable improvement of K availability to plants.

9.2.2 Potentials of Microbial Inoculants in Soil Structure Enhancement

Soil structure is a crucial aspect of sustainability in agriculture and ecosystem functioning because of its influence on the biological, physical and chemical properties of the soil. It refers to the three dimensional arrangement of organic or mineral complexes (aggregates) and pore spaces, which is usually quantified by size distribution of aggregate or the stability of aggregates. Aggregate formation and stabilization are mediated by several factors which include soil microorganisms (Lucas et al. 2013).

The role of microorganisms in the aggregate formation and stabilization of soil is well documented (Lucas et al. 2013; Helliwell et al. 2014). Activities of bacteria and fungi applied as inoculants in the enhancement of soil structure are affected by the aggregate scale (micro- or macroaggregate), soil types and soil mineralogy (Six et al. 2004). Aggregates are divided into microaggregate (<250 μm) and macroaggregate (>250 μm) and this division influences bacteria and fungi differently. While fungi stabilize macroaggregates, bacteria are more involved in the enhancement of microaggregates (Bossuyt et al. 2001). Bacteria play less role in coarse textured sandy soil where only the hyphal network is able to cross-link the abundant sand particles to form stable aggregates, whereas in clayey soil, both bacteria and fungi and their product play the role in aggregation (Six et al. 2004). Fungi are unique in influencing soil aggregate formation and stabilization because of the hyphae development and production of extra cellular polysaccharides. Hyphal networks enmesh and entangle macroaggregates while extracellular polysaccharides help to bind the micro-aggregates into stable macroaggregates (Bossuyt et al. 2001). Bacterial inoculants could play a key role in the soil structural stabilization through their secretions and exudates for microaggregate formation and stabilization.

The most studied fungi in soil structure stabilization are mycorrhizal fungi. Mycorrhizas are well recognized for their role in the improvement of soil structure (Leifheit et al. 2014). According to Rillig and Mummey (2006), mycorrhiza can influence soil aggregation at three main different scales; plant community, individual host plant root and fungal mycelium. Mycorrhizal ability to affect the plant community composition and cause root morphological changes to individual plants is well established (Oláh et al. 2005). However, our focus in this review is on the fungal mycelium which develops with the application of the fungal inoculant. Arbuscular mycorrhiza contribute to soil structure by 1) developing extraradical hyphae into the soil that align soil particles, providing the skeletal structure that enmeshes microaggregates to form macroaggregates; 2) secreting product like glomalin and glomalin related protein, mucilage, polysaccharides, hydrophobins and other extracellular compounds that cement aggregates and 3) delivering plant-derived carbon to aggregate surfaces (Rillig and Mummey 2006; Cardoso and Kuyper 2006). These processes are important for soil aggregation because of the space occupied by arbuscular mycorrhizal fungi in the soil system. Arbuscular mycorrhizal fungi produce significant biomass and represent dominant fungal biomass in agricultural soil (Rillig and Mummey 2006) and this is probably the reason for the positive effect of arbuscular mycorrhizal fungi inoculation on soil aggregation as reported by Leifheit et al. (2014). Considering the agricultural practices that are damaging to the soil structure, the use of mycorrhizal inoculants will not only help in the nutrition of crops, but also enhance the structural stability of agricultural soil. Both bacteria and fungi inoculants show potential for use in sustainable soil aggregate formation and stabilization and hence, soil structure enhancement.

9.2.3 *Role of Microbial Inoculants in Crop Tolerance to Drought Stress*

By reason of global climate change, drought is becoming more frequent and extreme in most part of the world. In most ecosystems, both fungi and bacteria are capable of resisting drought condition. However fungi show greater resistance than bacteria. Yuste et al. (2011), reported that fungi persisted longer in forest and desert soils during drought than bacteria. Arbuscular mycorrhizal fungi and saprophytic fungi have also exhibited better resistance to a wider range of heat and drought conditions compared to bacteria, However actinomycetes was an exception (Bell et al. 2009). This is not far from the fact fungi have extensive hyphal networks that enable them to access a larger volume of soil. These help fungi to regulate osmotic stress more effectively than bacteria (Leifheit et al. 2014). In the same vein Arbuscular mycorrhizal fungi show greater tolerance to drought than the saprophytic group (Davinic et al. 2013). This is associated with its ability to enhance greater plant nutrient and water uptake, greater carbon assimilation efficiencies. Fungi are also able to break-down more complex organic structures such as cellulose and lignin (Schwarze et al. 2004).

The discovery of some soil microorganism associated with natural drought condition in different ecological conditions, has necessitate their use as inoculants in drought season. The mechanisms by which these inoculants enhance plant drought tolerance include: increased hydric content, decreased antioxidant enzymatic activities, increased nutrient uptake, and decreased stomata conductance. They are also able to maintain indole acetic acid and increase proline production. Arbuscular mycorrhizal inoculants can improve crop drought tolerance in crop via glomalin induced changes in soil structure. Microbial inoculation during drought increased; plant growth, physiological and biochemical plant values that aid adaptive plant response, root growths, water content and plant C, K, Ca and Mg content (Armada et al. 2014). Some microorganisms that have been used to improve crop tolerance to drought include: Arbuscular mycorrhizal fungi such as *Glomus intraradices*, *Glomus mosseae*, *Aspergillus niger*, *Phanerochaete chrysosporium* (Medina et al. 2010; Wu et al. 2013), *Bacillus megaterium* (Armada et al. 2014), *Burkholderia phytofirmans* PsJN, *Enterobacter* sp. FD17 (Naveed et al. 2014), *Pseudomonas putida* (Armada et al. 2014), *Azospirillum* sp (Moutia et al. 2010), *Bacillus thuringiensis*, *Rhizophagus intraradices* (Ortiz et al. 2015).

9.2.4 Microbial Inoculants in the Remediation of Contaminated Soil

Bioaugmentation of tolerant crops with microbial inoculants can enhance plant establishment and growth under stress conditions including in the presence of soil contaminants. Phytoremediation of contaminated soil assisted by microbial inoculants enhance plant growth through: Production of plant growth hormones such as indole acetic acid and cytokinins, essential nutrients released by nitrogen fixers' siderophore producers and phosphorus solubilizer and suppression of the production of stress producing ethylene and hence have the potential to aid phytoremediation. Microbial inoculants are capable of remediating both organic and inorganic soil contaminants (Alori 2015). Some plants and associated microbial inoculants in phytoremediation of some soil inoculants are shown in Table 9.2.

Plants, in association with microbial inoculant, can remove or transform contaminants into harmless substance. Microbial populations through the release of chelating agents, acidification, phosphate solubilization and redox changes, affect heavy metal mobility and availability to the plant. The use of microbial inoculants in phytoremediation of polluted soil is cost efficient than alternative engineering-based solutions such as incineration, soil excavation, or land filling of the contaminated materials. Site use and remediation can occur simultaneously. It is an in situ approach, It treats the contamination in place so that large quantities of soil, sediment or water do not have to be pumped out or dug up of the ground for treatment. It is environmentally friendly, i.e, poses no health risk to neither plant, human nor animal. It enhances soil nutrient availability to the plants. Require less equipment

Table 9.2 Some microbial inoculants in phytoremediation of contaminated soil

Remediator (plant)	Microbial inoculants	Contaminant	References
<i>Withania somnifera</i>	<i>Staphylococcus cohnii</i> subsp <i>urealyticus</i>	Lindane	Abhilash et al. (2011)
<i>Lolium</i> sp and <i>Medicago sativa</i>	<i>Enterobacter ludwigii</i>	Hydrocarbon	Yousaf et al. (2011)
<i>Cytisus striatus</i>	<i>Rhodococcus erythropolis</i>	Hexachlorocyclohexane	Becerra-Castro et al. (2013)
<i>Brassica juncea</i>	<i>Azotobacter</i> sp + <i>Pseudomonas</i> sp	Cadmium	Panwar et al. (2011)
<i>Arabidopsis thaliana</i>	<i>Achromobacter xylosoxidans</i> F3B	Aromatic compounds	Ho et al. (2012)
<i>Lolium multiflorum</i>	<i>Pseudomonas</i> sp. ITRH76, <i>Rhodococcus</i> sp. ITRH43	Diesel	Afzal et al. (2011, 2012)
<i>Phragmites australis</i>	<i>Pseudomonas asplenii</i> AC	Copper, and creosote	Reed et al. (2005)
<i>Brassica juncea</i>	<i>Bacillus argabhattai</i> and <i>Bacillus megaterium</i>	Cadmium	Jeong et al. (2013)
<i>Lolium multiflorum</i>	<i>Pseudomonas putida</i> PCL1444	Naphthalene	Kuiper et al. (2004)
<i>Arabidopsis thaliana</i>	<i>Paenibacillus yonginensis</i>	Aluminium	Sukweenadhi et al. (2015)
<i>Lolium multiflorum</i>	<i>Pseudomonas nitroreducens</i> PS-2	Chlorpyrifos	Korade and Fulekar (2009)
<i>Phragmites australis</i>	<i>Sinorhizobium meliloti</i> P221	Phenanthrene	Golubev et al. (2009)
<i>Triticum aestivum</i>	<i>Azospirillum lipoferum</i>	Crude oil	Muratova et al. (2005)
<i>Vigna mungo</i>	<i>Pseudomonas aeruginosa</i> MKRh3	Cadmium	Ganesan (2008)
<i>Pisum sativum</i>	<i>Pseudomonas putida</i> VM1441 (pNAH7)	Naphthalene	Germaine et al. (2009)
<i>Phragmites australis</i>	Autochthonous microorganism consortium	Copper	Oliveira et al. (2014)
<i>Juncus maritimus</i> and <i>Phragmites australis</i>	Autochthonous microorganism consortium	Cadmium	Teixeira et al. (2014)

and labour than other methods, Phytoremediation using microbial inoculants does not degrade the physical or chemical health of the soil, unlike the soil excavation method that removes the topsoil that is rich in organic-matter- and the heavy machinery used compact the soil that is left behind. Microbial assisted phytoremediation do not require digging up or hauling of soil, hence it saves energy (Alori 2015). The strategies of microbial inoculants in remediation of polluted soil are safe and the effects are sustainable.

9.2.5 Benefits of Microbial Inoculation in Saline Soil

Salination of agricultural soil has become a serious threat to food production and security. According to (Shirmadi et al. 2010) about 5 % of the world soil is currently affected salinity. Vinocur and Altman (2005), predicted that by the year 2050, about 50 % of agricultural soils will be affected by salinity increase. Salinity has a direct effect on both the physical-chemical and biological properties of the soil, rendering such soils unsuitable for crop growth and biological processes. High soil salinity results in disruption in the uptake and transformation of nutrient elements such as Mg^{2+} and Ca^{2+} by plant. More also, it reduces ion activity in soil solution thereby, leading to nutrient deficiency and reduction of overall growth and yield quality of plant. Paul and Nair (2008), stress that plants become vulnerable to soil borne diseases under saline stress. In the past, some of the strategies employed to alleviate salt stress include the following: leaching of excess soluble salts from upper to lower soil depth, developing salt resistant cultivars, harvesting salt accumulating aerial plant parts in areas with negligible irrigation (Karthikeyan et al. 2012). These strategies are labour intensive and highly scientific. As a result, cost of cultivation may become increased and sometimes impossible. This has necessitated the need to discover agronomic system that can support plant growth under salinity stress that will not be accompanied by any environmental or health hazard.

Some soil microorganisms have been identified to be capable of alleviating salinity stress in plants and thus improving plants growth and yield. These soil microorganisms include the following genera: *Agrobacteria*, *Azospirillum*, *Bacillus*, *Glomus* *Gordonia* and *Pseudomonas*. They are environmentally-friendly, economically viable and energy efficient. The application of these groups of microorganisms is therefore a promising approach for alleviating salinity stress in plants.

Microbial inoculants ameliorate salt stress in plant via increased nutrient uptake, induced antioxidative defense system, modulation of the level of plant hormones, and reduction of ethylene level by producing 1-aminocyclopropane-1-carboxylate-deaminase in plants rhizosphere. Inoculation of sunflower with *Pseudomonas fluorescens* biotype F and *Pseudomonas fluorescens* CECT 378^T in sun flower grown in substrate with addition of salt (NaCl) showed that these strains that alleviate salt stress in sun flower produced indole-3-acetic acid and siderophores. The crop plants inoculated had a better developed root and a better K^+ : Na^+ ratio in the shoot (Shilev et al. 2012). In the same vein, Jha et al. (2011) discovered that the inoculation of a local paddy rice with *Pseudomonas pseudoacaligenes* and *Bacillus pumilus* in saline soil resulted in a decrease in growth suppression evident by an increased dry weight. The microbial inoculants also induced some osmoprotectants which help to overcome the deleterious effects of salt stress.

9.2.6 *Benefits of Microbial Inoculations as Biofertilizer on Plants Growth*

Some reports in literature collated recently by Babalola and Glick (2012) describe microbial inoculation to improve plant fitness and plant yield components. Microbial inoculation improves most plants growth and vigor. They enhance root growth and exudation (Babalola 2010; Trabelsi and Mhamdi 2013). When applied seeds, plants surface or soil, microbial inoculants increase the availability and supply of essential nutrients to host plants and thereby promoting growth. Microbial-inoculated plants show a reduction in membrane potential, accelerated osmotic adjustment, and enhanced lateral root development due to higher nitric acid and indole-3-acetic acid production (Dimkpa et al. 2009). Fungal inoculants will harmonize with the plant's root system and greatly expand the surface area of the root mass. Production of phytohormones by microbial inoculants can result in modification of root morphogenesis and hence support water uptake to plant roots.

Some common microbial components of biofertilizers include: *Azotobacter*, *Azospirillum*, *Bradyrhizobium*, mycorrhizae, phosphorus solubilizing bacteria, and *Rhizobium*. Microbial biofertilizers could be grouped into; Nitrogen fixers e.g. *Rhizobium* and *Bradyrhizobium*, phosphate solubilizers e.g. *Pseudomonas*, *Bacillus*, *Aspergillus* etc., cellulose degraders such as *Cytophaga* and phosphate mobilizers such as mycorrhizae. Microbial bio-fertilizers are cost effective and cheaper than the conventional techniques. They provide 25–30 % of chemical fertilizer equivalent of nitrogen. They increase phosphorus and potassium, increase water absorption and keep soil biologically active. In soils cropped with legumes, the application of arbuscular mycorrhizal fungi inoculants tremendously improve growth and yields. More also, inoculation with arbuscular mycorrhizal fungi improved growth of chickpea (*Cicer arietinum* L.) and doubled P uptake at low and intermediate levels of P in a pot experiment on sterilized low-P calcareous soil (Mohammadi et al. 2011). The inoculation of maize with *Trichoderma harzianum* strain T22 as a biofertilizer shortens the plant growth period and time and reduced lignifications hence, enhanced fresh state of maize plant (Akladios and Abbas 2012). An improved grain yield was reported by (N'Cho et al. 2013) when soybean was co-inoculated with rhizobium and fungal inoculants and application of foliar fertilizer.

However, microbial biofertilizers are associated with the following limitations:

- (i) The performance and efficacy of microbial inoculants (Biofertilizers) cannot be easily tested in the field i.e. there is a block in biofertilizer development.
- (ii) The efficacy of biofertilizers is not reliable. The mechanism of action of the biofertilizers in promoting growth is not yet well understood. In attempting to deal with these issues, research into biofertilizer is increasing.
- (iii) The essential nutrient may not be available in sufficient quantities to plants. Nutritional deficiency could exist due to low transfer of micro and macro nutrients

9.2.7 Benefits of Microbial Inoculants as Biocontrol Agents

Microbial inoculants have offered eco-friendly control mechanism against plant pathogens. Microbial inoculants produce antifungal secondary metabolites such as 2, 4-diacetylphloroglucinol and lytic enzymes. Some also confer plant protection against the activities of diseases causing organism by producing chitinase and protease enzymes. Microbial biocontrol agents also antagonize pathogens by competitive colonization of plant root and by forming biofilms in the hydroponic and soil systems.

Numerous microbial inoculants for control of several diseases especially species of the bacteria *Pseudomonas*, *Bacillus*, *Enterobacter*, *Streptomyces* and the fungus *Trichoderma*-, causing plant diseases as leaf spots, brown patch, *Pythium* blight and root rot, *Fusarium* wilt, dollar spot, summer patch, take-all patch, *Verticillium* wilt and *Typhula* blight have been studied by various researchers. Table 9.3 shows some micro-organisms that had exhibited some biocontrol activity against some phytopathogens.

Table 9.3 Microbial inoculants used as biocontrol agents

Biocontrol agents	Plant disease/Pathogen	Crop	References
<i>Bacillus subtilis</i> HJ5	<i>Verticillium</i> wilt	Cotton	Li et al. (2013)
<i>Bacillus subtilis</i> SQR9	<i>Fusarium</i> wilt	Cucumber	Cao et al. (2011)
<i>Bacillus subtilis</i>	<i>Fusarium</i> wilt	Maize	Cavaglieri et al. (2005)
<i>Trichoderma asperellum</i> T-34	<i>Rhizoctonia solani</i>	Cucumber	Trillas et al. (2006)
<i>Trichoderma harzianum</i> SQR-T 037	<i>Fusarium</i> wilt	Cucumber	Yang et al. (2011)
<i>Paenibacillus polymyxa</i> , <i>Trichoderma harzianum</i>	<i>Fusarium</i> wilt	Water melon	Wu et al. (2009)
<i>Streptomyces</i> sp strain g10	<i>Fusarium</i> wilt	Banana	Getha et al. (2005)
<i>Pseudomonas</i> spp	<i>Verticillium</i> wilt	Cotton	Erdogan and Benlioglu (2010)
<i>Bacillus pumilus</i> SQR-N43	<i>Rhizoctonia solani</i>	Cucumber	Huang et al. (2012)
<i>Streptomyces mutabilis</i> NBRC 12800	<i>Rhizoctonia solani</i> damping-off	Tomato	Goudjal et al. (2014)
<i>Bacillus amyloliquefaciens</i>	Panama disease	Banana	Xue et al. (2015)
<i>Streptomyces</i>	<i>Phytophthora</i> root rots	Alfalfa	Xiao et al. (2002)
<i>Streptomyces</i>	Damping-off	Sugar beet	Sadeghi et al. (2006)
<i>Actinoplanes campanulatus</i> , <i>Micromonospora chalcone</i> and <i>Streptomyces spiralis</i>	<i>Pythium aphanidermatum</i>	Cucumber	El-Tarabily et al. (2008)
<i>Streptomyces</i>	<i>Sclerotium rolfsii</i>	Sugar beet	Errakhi et al. (2007)
<i>Streptomyces</i>	Damping off	Tomato	Dhanasekaran et al. (2005)
<i>Micromonospora aurantiaca</i> , <i>Streptomyces griseus</i>	Damping off	Wheat	Hamdali et al. (2008)
<i>Bacillus pumilus</i> , <i>Pseudomonas alcaligenes</i> , and <i>Rhizobium</i> sp.	Wilt disease	Lentil	Akhtar et al. (2010)
<i>Bacillus subtilis</i>	Stem-end rot	Avocado flowers	Demoz and Korsten (2006)

9.3 Characteristics of Good Inoculants

Most microbial inoculants have received attention because of their catabolic versatility (Hmaeid et al. 2014), excellent root-colonizing ability which includes; motility, adhesion and growth rate (Hmaeid et al. 2014). They have capacity to produce a wide range of enzymes and metabolites, the ability to produce auxin or indole acetic acid, solubilize phosphate, produce siderophores (Hmaeid et al. 2014), survive and multiply in microhabitats associated with the root surface, in competition with other microbiota (Nivedhitha et al. 2008). They can persist in soil, are stable in storage and culture and are able to tolerate environmental constraints such as stress caused by fluctuating soil water conditions, use of fertilizers or agrochemicals (both organic and conventional) and soil disturbance such as cultivation (Hungaria et al. 2005). The success of microbial inoculation depends largely on the following: the plant species and cultivar, soil type, soil moisture and temperature conditions, the number of pathogens present in the soil around the plant and how the inoculants were prepared and applied (Babalola et al. 2007).

9.4 Properties of Good Carriers for Microbial Inoculants

The use of appropriate carriers for microbial inoculants preparation cannot be over emphasized. Good microbial inoculants carrier should (i) be easily handled and stored for a long period of time. (ii) have the capacity to deliver the right number of viable microbial cells in appropriate physiological condition at the right time, (iii) protect microbial cells from various biotic and abiotic stresses they will face once applied to the soil, (iv) retain microbial Plant-Growth Promoting abilities after a long period of storage, (v) be of low cost and locally available, (vi) be mixable and package able, (vii) permit gas exchange, particularly oxygen and have high organic matter content and water holding and retention capacity and it should be more than 50 %, (viii) be easy to process (mixing, curing and packaging operations) and free of lump- forming materials, (ix) be easy to sterilize by autoclaving or gamma-irradiation, (x) have good adhesion of seeds (xi) have good pH buffering capacity and (xii) be nontoxic to plants (Ferreira and Castro 2005).

Bacterial inoculants should be kept under a cool temperature, between 1.1 °C and 21.1 °C is best, away from extreme heat, direct sunlight or exposure to the elements or repeated freezing and thawing. Fungal inoculants are best kept dry. Excessive heat or cold is never of benefit. Also to be noted is the fact that agronomic practices have profound effects on soil organisms. They should therefore be designed to work in harmony with microbial inoculants and biological processes in order to support sustainable agricultural systems. Table 9.4 shows some materials that have successfully been used as carrier for microbial inoculants and the associated microbes. The use of appropriate carrier material determines the success of microbial inoculation techniques

Table 9.4 Types of carriers used for inoculants production

Carrier material	Inoculants	References
Sterilized oxalic acid, sludge, industrial waste, alginate-perlite dry granules, soybean oil or peanut oil added with lyophilized cells, composted sawdust, nutrient supplemented pumice, mineral soils, diatom, porosil mp, microcel, vermiculite, agriperlite, expanded clay, kaolin, celite, wheat bran, sugarcane bagasse, coal/charcoal, granular inoculants amended with nutrient and perlite	<i>Rhizobium</i> , <i>Sinorhizobium</i> , <i>Bradyrhizobium</i> , <i>Azospirillum</i> , <i>Agrobacterium</i> , Phosphorus solubilizing fungi, and <i>Aspergillus niger</i> ,	Zaidi et al. (2014)
Alginate beads supplemented with skim milk, charcoal based, broth based	<i>Bacillus subtilis</i> , <i>Pseudomonas corrugate</i>	Trivedi et al. (2012)
Talc powder +carboxyl- methyl cellulose	<i>Pseudomonas fluorescense</i>	Negi et al. (2005)

9.5 Methods of Application of Microbial Inoculants

The soil environmental conditions for crop production can be optimized by introducing friendly environmental microbial formulations to the soil. The use of appropriate carriers for microbial inoculants preparation is critical for the success of microbial inoculation (Babalola 2010). Microbial inoculant formulations are sold as wet able powders, granules or liquids sprays (Babalola and Glick 2012). The following methods can be used (Table 9.5). Microbial inoculants could be made of a singular strain of microbe. This approach is called the monoculture approach, where as an inoculant made of two or more strain of microbe or different types of organism is referred to as a co- culture or multiple culture approach. Microbial inoculants could either be applied directly to the soil or as seed/seedling dressing.

9.6 Factors that Determine the Performance of Microbial Inoculants Under Field Condition

The fate of introduced microbial inoculants includes: the ability to survive inoculation on seed, multiply in the spermosphere in response to seed exudates, attach to the root surface and colonize the developing root system. The ability of microbial inoculants to compete with indigenous microorganism present in the rhizosphere and the soil, for successful colonization of a developing plant depends on a number of biotic and abiotic factors. The survival, colonization and establishment of the inoculated microbes depend largely on these factors.

For microbial inoculants to survive competition, they must be able to sense chemo attractants like lipopolysaccharide such as O-antigen chain. However, lipopolysaccharide in colonization is strain dependent. O-antigenic side chain of *Pseudomonas fluorescens* PCL1205 is involve in tomato root colonization whereas

Table 9.5 Methods of application of microbial inoculants

Methods	Mechanism of application	Advantages and limitation	References
Directly to the soil	After seed germination, they are applied directly to the soil at the plant base near the plant roots	Withstand low moisture conditions better than carrier based inoculants. A less expensive method.	Mokone and Babalola (2013)
Seed application	Seeds are coated with microbe-carrier slurry. Adhesive solution such as sucrose solution is recommended.	Adequate loading of bacterial cells. Seeds treated with microbial inoculants may come in direct contact with any seed applied with chemicals which may adversely affect the survivability of the inoculated organism. Microbial culture may move away from rooting zones after application and could be exposed to agrochemicals after planting (Zaidi et al. 2014)	Mokone and Babalola (2013), Babalola et al. (2007), Babalola (2010) and Akladios and Abbas (2012)
Seedling root dip method	The seedling root dip method is mostly used for transplanted crops like vegetables. The roots of the seedling are dipped in a mixture of microbial culture and water for 5–10 min. The seedlings is then removed and transplanted almost immediately.	A less expensive method compare to carrier base inoculants	Babalola et al. (2007)
Field/Soil application	The direct application of inoculants to soil. Generally the granular inoculants are placed on the furrow under or alongside the seed. This enhanced inoculated microbe is in contact with the plant root.	less time consuming than the seed inoculation method	(Babalola et al. 2007)
Broadcasting method	Microbial inoculants could also be mixed with farmyard manure before broadcast.	Rapid and greater colonization of inoculants per unit area	(Akladios and Abbas 2012)

O-antigenic aspect of lipopolysaccharide of *Pseudomonas fluorescens* WCS374 does not contribute to rhizosphere colonization. Other factors include; high microbial growth rate and the ability of the inoculants to produce vitamin B1 and exude nicotinamide adenine dinucleotide dehydrogenases (NADH) and their ability to secrete site specific recombinase (Dennis et al. 2010).

The association between inoculated culture and host plant play a vital role in determining the success of microbial inoculation technique in the field. More also, Different inoculants produce different level and types of organic acid. The organic acids also vary in their ability to form complexes with cations to release inorganic nutrients for plant use. For instance, the ability of organic acid to complex with cation and liberate inorganic phosphorus varies with oxalic and citric acid. Gluconic acid has a limited ability to chelate and release phosphorus complex with calcium. Other factors include: Physico-chemical properties of soil such as soil pH, organic matter content and moisture content, presence of environmental pollutants such as xenobiotics and composition of root exudates. Further, management practices such as irrigating; grooming and fertilizing also influence microbial activity and growth. To overcome these short comings, applications must occur at times when environmental conditions strongly favor activity of the inoculant and inoculant must be formulated in a way that favors its activity and survival. The fate of microbial inoculants under field application depends largely on both biotic and abiotic factors.

9.7 Effects of Microbial Inoculants on the Resident Microbial Community

In the inoculation of seed and soil large quantity of efficient and viable microbial cells are introduced to the soil to cause a rapid colonization of the host rhizosphere. This may greatly disturb the equilibrium of soil microbial communities (Babalola 2014). These changes could either be by reason of direct trophic competitions or because of antagonistic or synergetic interactions between the introduced microbes and the resident microbes. It could also be indirect effect mediated by enhanced root growth and exudation. These changes could be in the taxonomic group or in the functional capabilities of the soil microbial community.

Depending on the technique used to address the effect of microbial inoculants on soil microbial communities, microbial inoculation may cause tremendous changes in the composition and number of taxonomic groups. While some researchers reported a long term effect, some other ones observed a transient or no effect at all. Plant and soil are affected by both the temporal and long term effects of inoculants. These effects result in unpredictable reactions (Trabelsi and Mhamdi 2013). For instance, Probanza et al. (2002) observed alteration in microbial rhizosphere composition when *Pinus pinea* L was inoculated with *Bacillus licheniformis* CECT 5106 and *Bacillus pumilus* CECT105. Conn and Franco (2004) observed that the introduction of a non-adapted (mixed commercial inoculants) microbial inoculums to the soil cropped with wheat disrupted the natural actinobacterial endophyte population thus reducing the diversity and colonization level. In contrast addition of a single actinobacterial endophyte to wheat plant increase colonization level and the indigenous endophyte population was not adversely affected (Yousaf et al. 2011). When maize was inoculated with *Azospirillum brasilense*, according to

Herschkovitz et al. (2005) the inoculants did not disrupt or alter the diversity and structure of root associated bacterial group both when universal bacterial primer and polymerase chain reaction (PCR)-denaturing gradient gel electrophoresis (DGGE) approach in conjunction with group-specific primers techniques were employed. The application some microbial inoculants could cause a change (which could be a decrease or an increase) in the equilibrium of soil microbial communities while some produce no effect at all.

9.8 Conclusion

This review has undoubtedly shown that microbial inoculants could improve biological management of nutrients and plant diseases resulting in improved plant performance in integrated plant management systems. They can contribute to a possible reduction of overuse of agro-chemicals and their environmental impacts. Though there are inconsistencies in the ability of many of the microorganisms that are used in the inoculant formulations to promote plant growth, the prospects of the inoculants like *rhizobium*, arbuscular mycorrhizal fungi and some *rhizobacteria* outweigh the lapses in meeting the goal of sustainable agricultural production. Apart from plant growth promotion, soil degradation is a serious problem in agriculture and the use of microbial inoculants is potentially part of the solution to this problem. More research effort is needed to elucidate the complex soil-plant-microbe interaction in order to reach the goal of completely substituting the environmentally degrading agro-chemicals with environmentally enhancing microbial inoculants. A combination of microorganisms in inoculant formulations has been shown to be helpful in many cases and can be made more efficient with research. Biotechnological research for effective and efficient microorganisms that are compatible with crops and adaptable to the soil environment will also be very helpful. In meeting the goal of sustainable agriculture, the use of microbial inoculants technology could be adopted for safe, increased production and sustainable agriculture.

Acknowledgements North-West University is gratefully acknowledged for ETA and MOD post-doctoral supports. OOB would like to thank the National Research Foundation, South Africa for grant (Ref: UID81192) that have supported research in her laboratory.

References

- Abhilash PC, Srivastava S, Srivastava P, Singh B, Jafri A, Singh N (2011) Influence of rhizospheric microbial inoculation and tolerant plant species on the rhizoremediation of lindane. *Environ Exp Bot* 74:127–130. doi:[10.1016/j.envexpbot.2011.05.009](https://doi.org/10.1016/j.envexpbot.2011.05.009)
- Afzal M, Yousaf S, Reichenauer TG, Kuffner M, Sessitsch A (2011) Soil type affects plant colonization, and catabolic gene expression of inoculated bacterial strains during phytoremediation of diesel. *J Hazard Mater* 186:1568–1575. doi:[10.1016/j.jhazmat.2010.12.040](https://doi.org/10.1016/j.jhazmat.2010.12.040)

- Afzal M, Yousaf S, Reichenauer TG, Sessitsch A (2012) The inoculation method affects colonization and performance of bacterial inoculant strains in the phytoremediation of soil contaminated with diesel oil. *Int J Phytoremediation* 14:35–47. doi:[10.1080/15226514.2011.552928](https://doi.org/10.1080/15226514.2011.552928)
- Ahemad M, Kibret M (2014) Mechanisms and applications of plant growth promoting rhizobacteria: current perspective. *J King Saud Univ Sci* 26(1):1–20. doi:[10.1016/j.jksus.2013.05.001](https://doi.org/10.1016/j.jksus.2013.05.001)
- Akhtar MS, Shakeel U, Siddiqui ZA (2010) Biocontrol of Fusarium wilt by *Bacillus pumilus*, *Pseudomonas alcaligenes*, and *Rhizobium* sp. on lentil. *Turk J Biology* 34:1–7. doi:[10.3906/biy-0809-12](https://doi.org/10.3906/biy-0809-12)
- Akladios SA, Abbas SA (2012) Application of *Trichoderma harziunum* T22 as a biofertilizer supporting maize growth. *Afr J Biotechnol* 11(35):8672–8683. doi:[10.5897/AJB11.4323](https://doi.org/10.5897/AJB11.4323)
- Alori ET (2015) Phytoremediation using microbial community II. In: Ansari AA, Gill SS, Newman L, Lanza GR (eds) *Phytoremediation: management of environmental contaminants*, vol II. Springer Publications, New York, pp. 183–190. doi:[10.1007/978-3-319-10969-5_15](https://doi.org/10.1007/978-3-319-10969-5_15)
- Alori E, Fawole O (2012) Phytoremediation of soils contaminated with aluminium and manganese by two arbuscular mycorrhizal fungi. *J Agric Sci* 4(8):246–252. doi:[10.5539/jas.v4n8p246](https://doi.org/10.5539/jas.v4n8p246)
- Ansari MF, Tipre DR, Dave SR (2014) Efficiency evaluation of commercial liquid biofertilizers for growth of *Cicer arietinum* (chickpea) in pot and field study. *Biocatal Agric Biotechnol* 4(1):17–24. doi:[10.1016/j.bcab.2014.09.010](https://doi.org/10.1016/j.bcab.2014.09.010)
- Armada E, Portela G, Roldán A, Azcóna R (2014) Combined use of beneficial soil microorganism and agrowaste residue to cope with plant water limitation under semiarid conditions. *Geoderma* 232–234:640–648. doi:[10.1016/j.geoderma.2014.06.025](https://doi.org/10.1016/j.geoderma.2014.06.025)
- Babalola OO (2010) Beneficial bacteria of agricultural importance. *Biotechnol Lett* 32(11):1559–1570. doi:[10.1007/s10529-010-0347-0](https://doi.org/10.1007/s10529-010-0347-0)
- Babalola OO (2014) Does nature make provision for backups in the modification of bacterial community structures? *Biotechnol Genet Eng Rev* 30(1):31–48. doi:[10.1080/02648725.2014.921497](https://doi.org/10.1080/02648725.2014.921497)
- Babalola OO, Glick BR (2012) Indigenous African agriculture and plant associated microbes: current practice and future transgenic prospects. *Sci Res Essays* 7(28):2431–2439. doi:[10.5897/SRE11.1714](https://doi.org/10.5897/SRE11.1714)
- Babalola OO, Sanni AI, Odhiambo GD, Torto B (2007) Plant growth-promoting rhizobacteria do not pose any deleterious effect on cowpea and detectable amounts of ethylene are produced. *World J Microbiol Biotechnol* 23(6):747–752. doi:[10.1007/s11274-006-9290-6](https://doi.org/10.1007/s11274-006-9290-6)
- Barac T, Weyens N, Oeyen L, Taghavi S, van der Lelie D, Dubin D, Spliet M, Vangronveld J (2009) Field note: hydraulic containment of BTEX plume using poplar trees. *Int J Phytoremediation* 11:416–424. doi:[10.1080/15226510802655880](https://doi.org/10.1080/15226510802655880)
- Bargaz A, Faghire M, Abdi N, Farissi M, Sifi B, Drevon J-J, Ikbali MC, Ghoulam C (2012) Low soil phosphorus availability increases acid phosphatases activities and affects P partitioning in nodules, Seeds and Rhizosphere of *Phaseolus vulgaris*. *Agriculture* 2:139–153. doi:[10.3390/agriculture2020139](https://doi.org/10.3390/agriculture2020139)
- Becerra-Castro C, Prieto-Fernández Á, Kidd P, Weyens N, Rodríguez-Garrido B, Touceda-González M, Acea MJ, Vangronsveld J (2013) Improving performance of *Cytisus striatus* on substrates contaminated with hexachlorocyclohexane (HCH) isomers using bacterial inoculants: developing a phytoremediation strategy. *Plant Soil* 362:247–260. doi:[10.1007/s11104-012-1276-6](https://doi.org/10.1007/s11104-012-1276-6)
- Bell CW, Acosta-Martínez V, McIntyre NE, Cox S, Tissue DT, Zak JC (2009) Linking microbial community structure and function to seasonal differences in soil moisture and temperature in a Chihuahuan desert grassland. *Microb Ecol* 58:827–842. doi:[10.1007/s00248-009-9529-5](https://doi.org/10.1007/s00248-009-9529-5)
- Bernhard A (2010) The nitrogen cycle: processes, players, and human impact. *Nat Educ Knowl* 3(10):25
- Bossuyt H, Denef K, Six J, Frey SD, Merckx R, Paustian K (2001) Influence of microbial populations and residue quality on aggregate stability. *Appl Soil Ecol* 16:195–208. doi:[10.1016/S0929-1393\(00\)00116-5](https://doi.org/10.1016/S0929-1393(00)00116-5)

- Calvo P, Nelson L, Kloepper JW (2014) Agricultural uses of plant biostimulants. *Plant Soil* 383(1-2):3–41. doi:[10.1007/s11104-014-2131-8](https://doi.org/10.1007/s11104-014-2131-8)
- Cao Y, Ling N, Yang XM, Chen LH, Shen QR (2011) *Bacillus subtilis* SQR9 can control *Fusarium* wilt in cucumber by colonizing plant roots. *Biol Fertil Soils* 47:495–506. doi:[10.1007/s00374-011-0556-2](https://doi.org/10.1007/s00374-011-0556-2)
- Cardoso IM, Kuyper TW (2006) Mycorrhizas and tropical soil fertility. *Agric Ecosyst Environ* 116:72–84. doi:[10.1016/j.agee.2006.03.011](https://doi.org/10.1016/j.agee.2006.03.011)
- Carvajal-Muñoz JS, Carmona-García CE (2012) Benefits and limitations of biofertilization in agricultural practices. *Livest Res Rural Dev* 24(3)
- Cavaglieri L, Orlando J, Rodríguez MI, Chulze S, Etcheverry M (2005) Biocontrol of *Bacillus subtilis* against *Fusarium verticillioides* in vitro at the maize root level. *Res Microbiol* 156:748–754. doi:[10.1016/j.resmic.2005.03.001](https://doi.org/10.1016/j.resmic.2005.03.001)
- Chatzipavlidis I, Kefalogianni I, Venieraki A, Holzapfel W (2013) Status and trends of the conservation and sustainable use of microorganisms in agroindustrial processes. Food and Agricultural Organization (FAO) United Nations, United Nations
- Conn VM, Franco CMM (2004) Effect of Microbial inoculants on the indigenous Actinobacterial Endophyte Population in the Roots of Wheat as Determined by Terminal Restriction Fragment Length Polymorphism. *Appl Environ Microbiol* 70(11):6407–6413. doi:[10.1128/AEM.70.11.6407-6413.2004](https://doi.org/10.1128/AEM.70.11.6407-6413.2004)
- Cozzolino V, Di Meo V, Piccolo A (2013) Impact of arbuscular mycorrhizal fungi application on maize production and soil phosphorus availability. *J Geochem Explor* 129:40–44. doi:[10.1016/j.geexplo.2013.02.006](https://doi.org/10.1016/j.geexplo.2013.02.006)
- Dare MO, Abaidoo RC, Fagbola O, Asiedu R (2010) Effects of arbuscular mycorrhizal inoculation and phosphorus application on yield and nutrient uptake of yam. *Commun Soil Sci Plant* 41:2729–2743. doi:[10.1080/00103624.2010.518264](https://doi.org/10.1080/00103624.2010.518264)
- Davinic M, Moore-Kucera J, Acosta-Martínez V, Zak J, Allen V (2013) Soil fungal distribution and functionality as affected by grazing and vegetation components of integrated crop–livestock agroecosystems. *Appl Soil Ecol* 66:61–70. doi:[10.1016/j.apsoil.2013.01.013](https://doi.org/10.1016/j.apsoil.2013.01.013)
- Dennis PG, Miller AJ, Hirsch PR (2010) Are root exudates more important than other sources of rhizodeposits in structuring rhizosphere bacterial communities? *FEMS Microbiol Ecol* 72:313–327. doi:[10.1111/j.1574-6941.2010.00860.x](https://doi.org/10.1111/j.1574-6941.2010.00860.x)
- Demoz BT, Korsten L (2006) *Bacillus subtilis* attachment, colonization, and survival on avocado flowers and its mode of action on stem-end rot pathogens. *Biol Control* 37:68–74. doi:[10.1016/j.Biocontrol.2005.11.010](https://doi.org/10.1016/j.Biocontrol.2005.11.010)
- Dhanasekaran D, Sivamani P, Panneerselvam A, Thajuddin N, Rajakumar G, Selva-mani S (2005) Biological control of tomato seedling damping off with *Streptomyces* sp. *Plant Pathol J* 4:91–95
- Dimkpa C, Weinand T, Asch F (2009) Plant–rhizobacteria interactions alleviate abiotic stress conditions. *Plant Cell Environ* 321:682–1694. doi:[10.1111/j.1365-3040.2009.02028.x](https://doi.org/10.1111/j.1365-3040.2009.02028.x)
- Egamberdiyeva D (2007) The effect of plant growth promoting bacteria on growth and nutrient uptake of maize in two different soils. *Appl Soil Ecol* 36:184–189. doi:[10.1016/j.apsoil.2007.02.005](https://doi.org/10.1016/j.apsoil.2007.02.005)
- El-Tarabily KA, Nasser AH, Hardy GE, Sivaithamparam K (2008) Plant growth promotion and biological control of *Pythium aphanidermatum*, a pathogen of cucumber, by endophytic actinomycetes. *J Appl Microbiol* 106:13–26. doi:[10.1111/j.1365-2672.2008.03926.x](https://doi.org/10.1111/j.1365-2672.2008.03926.x)
- Erdoğan O, Benlioğlu K (2010) Biological control of *Verticillium* wilt on cotton by the use of fluorescent *Pseudomonas* spp. under field conditions. *BioControl* 53:39–45. doi:[10.1016/J.Biocontrol.2009.11.011](https://doi.org/10.1016/J.Biocontrol.2009.11.011)
- Errakhi R, Bouton F, Lebrihi A, Barakate M (2007) Evidence of biological control capacities of *Streptomyces* spp. against *Sclerotium rolfsii* responsible for damping-off disease in sugar beet (*Beta vulgaris* L.). *World J Microbiol Biotechnol* 23:1503–1509
- Ferreira EM, Castro IV (2005) Residues of the cork industry as carriers for the production of legumes inoculants. *Silva Lusitana* 13(2):159–167

- Gaind S (2011) Microbial inoculants: an approach to sustainable agriculture. Biotech Article
- Ganesan V (2008) Rhizoremediation of cadmium soil using a cadmium-resistant plant growth-promoting rhizopseudomonad. *Curr Microbiol* 56:403–407. doi:[10.1007/s00284-008-9099-7](https://doi.org/10.1007/s00284-008-9099-7)
- Germaine KJ, Keogh E, Ryan D, Dowling DN (2009) Bacterial endophyte-mediated naphthalene phytoprotection and phytoremediation. *FEMS Microbiol Lett* 296:226–234. doi:[10.1111/j.1574-6968.2009.01637.x](https://doi.org/10.1111/j.1574-6968.2009.01637.x)
- Getha K, Vikineswary S, Wong WH, Seki T, Ward A, Goodfellow M (2005) Evaluation of *Streptomyces* sp. strain g10 for suppression of *Fusarium* wilt, rhizosphere colonization in pot-grown banana plantlet. *J Ind Microbiol Biotechnol* 32:24–32. doi:[10.1007/s10295-004-0199-5](https://doi.org/10.1007/s10295-004-0199-5)
- Golubev S, Schelud'ko A, Muratova A, Makarov O, Turkovskaya O (2009) Assessing the potential of rhizobacteria to survive under phenanthrene pollution. *Water Air Soil Pollut* 198:5–16. doi:[10.1007/s11270-008-9821-x](https://doi.org/10.1007/s11270-008-9821-x)
- Goudjal Y, Toumatia O, Yekkour A, Sabaou N, Mathieu F, Zitouni A (2014) Biocontrol of *Rhizoctonia solani* damping-off and promotion of tomato plant growth by endophytic actinomycetes isolated from native plants of Algerian Sahara. *Microbiol Res* 169:59–65. doi:[10.1016/j.micres.2013.06.014](https://doi.org/10.1016/j.micres.2013.06.014)
- Halpern M, Bar-Tal A, Ofek M, Minz D, Muller T, Yermiyahu U (2015) The use of biostimulants for enhancing nutrient uptake. In: LS D (ed) *Advances in Agronomy*, vol 130. Academic Press, pp. 141–174. doi:[10.1016/bs.agron.2014.10.001](https://doi.org/10.1016/bs.agron.2014.10.001)
- Hamdali H, Hafidi M, Virolle MJ, Ouhdouch Y (2008) Growth promotion and protection against damping-off of wheat by two rock phosphate solubilizing actinomycetes in a P-deficient soil under greenhouse conditions. *Appl Soil Ecol* 40:510–517. doi:[10.1016/j.apsoil.2008.08.001](https://doi.org/10.1016/j.apsoil.2008.08.001)
- Helliwell JR, Miller AJ, Whalley WR, Mooney SJ, Sturrock CJ (2014) Quantifying the impact of microbes on structural development and behaviour in wet soil. *Soil Biol Biochem* 74:138–147. doi:[10.1016/j.soilbio.2014.03.009](https://doi.org/10.1016/j.soilbio.2014.03.009)
- Herschkovitz Y, Lerner A, Davidov Y, Okon Y, Jurkevitch E (2005) *Azospirillum brasilense* does not affect population structure of specific rhizobacterial communities of inoculated maize (*Zea mays*). *Environ Microbiol* 7(11):1847–1852. doi:[10.1111/j.1462-2920.2005.00926.x](https://doi.org/10.1111/j.1462-2920.2005.00926.x)
- Hmaeid N, Metoui O, Wali M, Zorrig W, Abdelly C (2014) Comparative effects of *Rhizobacteria* in promoting growth of *Hordeum maritimum* L. plants under salt stress. *J Plant Biol Res* 3(1):37–50
- Ho Y-N, Mathew DC, Hsiaoa S-C, Chun-Hao Shiha C-H, Chienb M-F, Chiang H-M, Huang C-C (2012) Selection and application of endophytic bacterium *Achromobacter xylosoxidans* strain F3B for improving phytoremediation of phenolic pollutants. *J Hazard Mater*. doi:[10.1016/j.jhazmat.2012.03.035](https://doi.org/10.1016/j.jhazmat.2012.03.035)
- Hodge A, Storer K (2015) Arbuscular mycorrhiza and nitrogen: implications for individual plants through to ecosystems. *Plant Soil* 383(1-2):1–19. doi:[10.1007/s11104-014-2162-1](https://doi.org/10.1007/s11104-014-2162-1)
- Hong S, Kim D, Baek S, Kwon S, Samson RA (2011) Taxonomy of *Eurotium* species isolated from meju. *J Microbiol* 49:669–674. doi:[10.1007/s12275-011-0376-y](https://doi.org/10.1007/s12275-011-0376-y)
- Huang X, Zhang N, Yong X, Yang X, Shen Q (2012) Biocontrol of *Rhizoctonia solani* damping-off disease in cucumber with *Bacillus pumilus* SQR-N43. *Microbiol Res* 167:135–143. doi:[10.1016/J.Micres.2011.06.002](https://doi.org/10.1016/J.Micres.2011.06.002)
- Hungaria M, Loureiro MF, Mendes IC, Campo RJ, Graham PH (2005) Inoculant preparation, production and application. In: Werner D, Newton WE (eds) *Nitrogen fixation in agriculture, forestry, ecology and environment*, vol 4. Springer, Netherlands, pp. 223–253. doi:[10.1007/1-4020-3544-6_11](https://doi.org/10.1007/1-4020-3544-6_11)
- Jenkinson DA (2001) The impact of human on the nitrogen cycle with focus on temperate arable agriculture. *Plant Soil* 228:3–15. doi:[10.1023/A:1004870606003](https://doi.org/10.1023/A:1004870606003)
- Jeong S, Moon HS, Shin D, Nam K (2013) Survival of introduced phosphate-solubilizing bacteria (PSB) and their impact on microbial community structure during the phytoextraction of Cd-contaminated soil. *J Hazard Mater* 263(2):441–449. doi:[10.1016/j.jhazmat.2013.09.062](https://doi.org/10.1016/j.jhazmat.2013.09.062)

- Jha Y, Subramanian RB, Patel S (2011) Combination of endophytic and rhizospheric plant growth promoting rhizobacteria in *Oryza sativa* shows higher accumulation of osmoprotectant against saline stress. *Acta Physiol Plant* 33:797–802. doi:[10.1007/s11738-010-0604-9](https://doi.org/10.1007/s11738-010-0604-9)
- Julia WG, Peter H, Elizabeth L, Glen H (2013) Soil inoculants. University of Georgia College of Agricultural and Environmental Sciences, Cooperative Extension 10 pages
- Karthikeyan B, Joe MM, Islam MR, Sa T (2012) ACC deaminase containing diazotrophic endophytic bacteria ameliorate salt stress in *Catharanthus roseus* through reduced ethylene levels and induction of antioxidative defense systems. *Symbiosis* 56:77–86. doi:[10.1007/s13199-012-0162-6](https://doi.org/10.1007/s13199-012-0162-6)
- Kavino M, Harish S, Kumar N, Saravanakumar D, Samiyappan R (2010) Effect of chitinolytic PGPR on growth, yield and physiological attributes of banana (*musa spp.*) under field conditions. *Appl Soil Ecol* 45:71–77. doi:[10.1016/j.apsoil.2010.02.003](https://doi.org/10.1016/j.apsoil.2010.02.003)
- Korade DL, Fulekar MH (2009) Rhizosphere remediation of chlorpyrifos in mycorrhizospheric soil using ryegrass. *J Hazard Mater* 172:1344–1350. doi:[10.1016/J.Jhazmat.2009.08.002](https://doi.org/10.1016/J.Jhazmat.2009.08.002)
- Kruger M, Kruger C, Walker C, Stockinger H, Schubler A (2012) Phylogenetic reference data for systematics and phylotaxonomy of arbuscular mycorrhizal fungi from phylum to species level. *New Phytol* 193:970–984. doi:[10.1111/j.1469-8137.2011.03962.x](https://doi.org/10.1111/j.1469-8137.2011.03962.x)
- Kuiper I, Lagendijk EL, Bloemberg GV, Lugtenberg BJJ (2004) Rhizoremediation: a beneficial plant-microbe interaction. *Mol Plant Microbe Interact* 17:6–15. doi:[10.1094/MPMI.2004.17.1.6](https://doi.org/10.1094/MPMI.2004.17.1.6)
- Kumar B, Trivedi P, Pandey A (2007) *Pseudomonas corrugata*: a suitable bacterial inoculant for maize grown under rainfed conditions of Himalayan region. *Soil Biol Biochem* 39(12):3093–3100. doi:[10.1016/j.soilbio.2007.07.003](https://doi.org/10.1016/j.soilbio.2007.07.003)
- Leifheit EF, Veresoglou SD, Lehmann A, Morris EK, Rillig MC (2014) Multiple factors influence the role of arbuscular mycorrhizal fungi in soil aggregation—a meta-analysis. *Plant Soil* 374:523–537. doi:[10.1007/s11104-013-1899-2](https://doi.org/10.1007/s11104-013-1899-2)
- Li Y, Zou YN, Wu QS (2013) Effects of inoculation with *Diversispora spurca* on growth, root system architecture and chlorophyll contents of four citrus genotype plants. *Int J Agric Biol* 15:342–346
- Liu D, Lian B, Dong H (2012) Isolation of *Paenibacillus* sp. and assessment of its potential for enhancing mineral weathering. *Geomicrobiol J* 29(5):413–421. doi:[10.1080/01490451.2011.576602](https://doi.org/10.1080/01490451.2011.576602)
- Lucas ST, D'Angelo EM, Williams MA (2013) Improving soil structure by promoting fungal abundance with organic soil amendments. *Appl Soil Ecol* 75:13–23. doi:[10.1016/j.apsoil.2013.10.002](https://doi.org/10.1016/j.apsoil.2013.10.002)
- Ma Y, Rajkumar M, Freitas H (2009) Inoculation of plant growth promoting bacterium *Achromobacter xylosoxidans* strain Ax10 for the improvement of copper phytoextraction by *Brassica juncea*. *J Environ Manag* 90:831–837. doi:[10.1016/j.jenvman.2008.01.014](https://doi.org/10.1016/j.jenvman.2008.01.014)
- Mokone PH, Babalola OO (2013) Evaluation of plant growth promoting potential of four rhizobacterial species for indigenous system. *J Cent South Univ* 20:164–171. doi:[10.1007/s11771-013-1472-4](https://doi.org/10.1007/s11771-013-1472-4)
- Medina A, Roldán A, Azcón R (2010) The effectiveness of arbuscular-mycorrhizal fungi and *Aspergillus niger* or *Phanerochaete chrysosporium* treated organic amendments from olive residues upon plant growth in a semi-arid degraded soil. *J Environ Manag* 91:2547–2553. doi:[10.1016/j.jenvman.2010.07.008](https://doi.org/10.1016/j.jenvman.2010.07.008)
- Meena VS, Maurya BR, Verma JP (2014) Does a rhizospheric microorganism enhance K⁺ availability in agricultural soils? *Microbiol Res* 169 (5–6):337–347. doi:<http://dx.doi.org/10.1016/j.micres.2013.09.003>
- Mohammadi K, Khalesro S, Sohrabi Y, Heidari G (2011) A review: beneficial effects of the mycorrhizal fungi for plant growth. *J Appl Environ Biol Sci* 1(9):310–319
- Moutia J-FY, Saumtally S, Spaepen S, Vanderleyden J (2010) Plant growth pro-motion by *Azospirillum* sp. in sugarcane is influenced by genotype and droughtstress. *Plant Soil* 337:233–242. doi:[10.1007/s11104-010-0519-7](https://doi.org/10.1007/s11104-010-0519-7)

- Muratova AY, Turkovskaya OV, Antonyuk LP, Makarov OE, Pozdnyakova LI, Ignatov V (2005) Oil-oxidizing potential of associative rhizobacteria of the genus *Azospirillum*. *Microbiology* 74:210–215
- Naveed M, Mitter B, Reichenauer TG, Wieczorek K, Sessitscha A (2014) Increased drought stress resilience of maize through endophytic colonization by *Burkholderia phytofirmans* PsJN and *Enterobacter* sp. FD17. *Environ Exp Bot* 97:30–39. doi:[10.1016/j.envexpbot.2013.09.014](https://doi.org/10.1016/j.envexpbot.2013.09.014)
- N'Cho CO, Yusuf AA, Ama-Abina JT, Jemo M, Abaidoo RC, Savane I (2013) Effects of commercial microbial inoculants and foliar fertilizers on soybean nodulation and yield in northern Guinea savannah of Nigeria. *Int J Adv Agric Res* 1:66–73
- Negi PS, Chauhan AS, Sadia GA, Rohinishree YS, Ramteke RS (2005) Antioxidant and antibacterial activities of various seabuckthorn (*hippophae rhamnoides* L.) seed extracts. *Food Chem* 92:119–124. doi:[10.1016/j.foodchem.2004.07.009](https://doi.org/10.1016/j.foodchem.2004.07.009)
- Nivedhitha VR, Shwetha B, Deepa DD, Manojkumar NH, Raghavendra RB (2008) Plant growth promoting microorganisms (PGPMs) from bamboo rhizosphere. *Adv Biotechnol*:33–35
- Oláh B, Brière C, Bécard G, Dénarié J, Gough C (2005) Nod factors and a diffusible factor from arbuscular mycorrhizal fungi stimulate lateral root formation in *Medicago truncatula* via the DMI1/DMI2 signalling pathway. *Plant J* 44:195–207
- Oliveira T, Mucha AP, Reis I, Rodrigues P, Gomes CR, Almeida CMR (2014) Copper phytoremediation by a salt marsh plant (*Phragmites australis*) enhanced by autochthonous bioaugmentation. *Mar Pollut Bull* 88:231–238. doi:[10.1016/j.marpolbul.2014.08.038](https://doi.org/10.1016/j.marpolbul.2014.08.038)
- Ortiz N, Armada E, Duque E, Roldán A, Azcón R (2015) Contribution of arbuscular mycorrhizal fungi and/or bacteria to enhancing plant drought tolerance under natural soil conditions: Effectiveness of autochthonous or allochthonous strains. *J Plant Physiol* 174:87–96. doi:[10.1016/j.jplph.2014.08.019](https://doi.org/10.1016/j.jplph.2014.08.019)
- Owen D, Williams AP, Griffith GW, Withers PJA (2015) Use of commercial bio-inoculants to increase agricultural production through improved phosphorous acquisition. *Appl Soil Ecol* 86:41–54. doi:[10.1016/j.apsoil.2014.09.012](https://doi.org/10.1016/j.apsoil.2014.09.012)
- Panwar BS, Kádár I, Bíró B, Rajkai-végh K, Ragályi P, Rékási M, Márton L (2011) Phytoremediation: Enhanced cadmium (Cd) accumulation by organic manuring, Edta and microbial inoculants (*Azotobacter* sp., *Pseudomonas* sp.) in Indian mustard *Brassica juncea* L). *Act Agron Hung* 59(2):117–123. doi:[10.1556/AAgr.59.2011.2.2](https://doi.org/10.1556/AAgr.59.2011.2.2)
- Parmar P, Sindhu SS (2013) Potassium solubilization by rhizosphere bacteria: influence of nutritional and environmental conditions. *J Microbiol Res* 3(1):25–31. doi:[10.5923/j.microbiology.20130301.04](https://doi.org/10.5923/j.microbiology.20130301.04)
- Porras-Soriano A, Soriano-Martin ML, Porras-Piedra A, Azcón R (2009) Arbuscular mycorrhizal fungi increased growth, nutrient uptake and tolerance to salinity in olive trees under nursery conditions. *J Plant Physiol* 166:1359. doi:[10.1128/AEM.71.12.8500-8505.2005](https://doi.org/10.1128/AEM.71.12.8500-8505.2005)
- Paul D, Nair S (2008) Stress adaptations in a plant growth promoting *Rhizobacterium* (PGPR) with increasing salinity in the coastal agricultural soils. *J Basic Microbiol* 48:1–7. doi:[10.1002/jobm.200700365](https://doi.org/10.1002/jobm.200700365)
- Pereg L, McMillan M (2015) Scoping the potential uses of beneficial microorganisms for increasing productivity in cotton cropping systems. *Soil Biol Biochem* 80:349–358. doi:[10.1016/j.soilbio.2014.10.020](https://doi.org/10.1016/j.soilbio.2014.10.020)
- Probanza A, García JAL, Palomino MR, Ramos B, Mañero FJG (2002) *Pinus pinea* L. seedling growth and bacterial rhizosphere structure after inoculation with PGPR *Bacillus* (*B. licheniformis* CECT 5106 and *B. pumilus* CECT 5105). *Appl Soil Ecol* 20:75–84. doi:[10.1016/S0929-1393\(02\)00007-0](https://doi.org/10.1016/S0929-1393(02)00007-0)
- Rawat AK, Rao DLN, Sahu RK (2013) Effect of soybean inoculation with *Bradyrhizobium* and wheat inoculation with *Azotobacter* on their productivity and N turnover in a vertisol. *Arch Agron Soil Sci* 59(11):1559–1571. doi:[10.1080/03650340.2012.740555](https://doi.org/10.1080/03650340.2012.740555)
- Reed MLE, Warner BG, Glick BR (2005) Plant growth-promoting bacteria facilitate the growth of the common reed *Phragmites australis* in the presence of copper or polycyclic aromatic hydrocarbons. *Curr Microbiol* 51(6):425–429. doi:[10.1007/s00284-005-4584-8](https://doi.org/10.1007/s00284-005-4584-8)

- Rillig MC, Mummey DL (2006) Mycorrhizas and soil structure. *New Phytol* 171:41–53. doi:[10.1111/j.1469-8137.2006.01750.x](https://doi.org/10.1111/j.1469-8137.2006.01750.x)
- Sadeghi A, Hessian AR, Askari H, Aghighi S, Shahidi BGH (2006) Biological control potential of two *Streptomyces* isolates on *Rhizoctonia solani*, the causal agent of damping-off of sugar beet. *Pak J Biol Sci* 9:904–910
- Saharan BS, Nehra V (2011) Plant growth promoting rhizobacteria: a critical review. *Life Sci Med Res* 21:1–30
- Sanderson RT (2014) Phosphorus (P) Chemical element. *Encyclopedia Britannica*
- Sangeeth KP, Bhai RS, Srinivasan V (2012) *Paenibacillus glucanolyticus*, a promising potassium solubilizing bacterium isolated from black pepper (*Piper nigrum* L.) rhizosphere. *J Spices Aromat Crops* 21(2):118–124
- Schwarze FWMR, Engels J, Mattheck C (2004) Fungal strategies of wood decay in trees. Springer-Verlag, Berlin
- Shilev S, Sancho ED, Benlloch-González M (2012) Rhizospheric bacteria alleviate salt-produced stress in sunflower. *J Environ Manag* 95(Supplement):S37–S41. doi:[10.1016/j.jenvman.2010.07.019](https://doi.org/10.1016/j.jenvman.2010.07.019)
- Shirmadi M, Savaghebi GR, Khavazi K, Akbarzadeh A, Farahbakhsh M, Rejali F, Sadat A (2010) Effect of microbial inoculants on uptake of nutrient elements in two cultivars of sunflower (*Helianthus annuus* L.) in saline soils. *Not Sci Biol* 2(3):57–66
- Singh B, Satyanarayana T (2011) Microbial phytases in phosphorus acquisition and plant growth promotion. *Physiol Mol Biol Plants* 17:93–103. doi:[10.1007/s12298-011-0062-x](https://doi.org/10.1007/s12298-011-0062-x)
- Singh R, Soni SK, Patel RP, Kalra A (2013) Technology for improving essential oil yield of *Ocimum basilicum* L. (sweet basil) by application of bioinoculant colonized seeds under organic field conditions. *Ind Crop Prod* 45:335–342. doi:[10.1016/j.indcrop.2013.01.003](https://doi.org/10.1016/j.indcrop.2013.01.003)
- Six J, Bossuyt H, Degryze S, Denef K (2004) Mycorrhizal Symbiosis. *Soil Tillage Res* 79:7–31
- Smith SE, Read DJ (2008) Mycorrhizal Symbiosis, 3rd edn. Academic Press, London
- Smith SE, Jakobsen I, Grønlund M, Smith FA (2011) Roles of arbuscular mycorrhizas in plant phosphorus nutrition: interactions between pathways of phosphorus uptake in arbuscular mycorrhizal roots have important implications for understanding and manipulating plant phosphorus acquisition. *Plant Physiol* 156:1050–1057. doi:[10.1104/pp.111.174581](https://doi.org/10.1104/pp.111.174581)
- Trillas MI, Casanova E, Corxarrera L, Ordovas J, Borrero C, Aviles M (2006) Composts from agricultural waste, the *Trichoderma asper-ellum* strain T-34 suppress *Rhizoctonia solani* in cucumber seedlings. *BioControl* 39:32–38. doi:[10.1016/j.biocontrol.2006.05.007](https://doi.org/10.1016/j.biocontrol.2006.05.007)
- Sukweenadhia J, Kima Y, Choib E, Kohc S, Leed S, Kima Y, Yanga DC (2015) *Paenibacillus yonginensis* DCY84 induces changes in *Arabidopsis thaliana* gene expression against aluminum, drought, and salt stress. *Microbiol Res* 172:7–15. doi:[10.1016/j.micres.2015.01.007](https://doi.org/10.1016/j.micres.2015.01.007)
- Taghavi S, Barac T, Greenberg B, Borremans B, Vangronsveld J, van der Lelie D (2005) Horizontal gene transfer to endogenous endophytic bacteria from poplar improves phytoremediation of toluene. *Appl Environ Microbiol* 71:8500–8505
- Tank N, Saraf M (2010) Salinity-resistant plant growth promoting rhizobacteria ameliorates sodium chloride stress on tomato plants. *J Plant Interact* 5:51–58. doi:[10.1080/17429140903125848](https://doi.org/10.1080/17429140903125848)
- Teixeira C, Almeida MR, da Silva MN, Bordalo AA, Mucha AP (2014) Development of autochthonous microbial consortia for enhanced phytoremediation of salt-marsh sediments contaminated with cadmium. *Sci Total Environ* 493:757–765. doi:[10.1016/j.scitotenv.2014.06.040](https://doi.org/10.1016/j.scitotenv.2014.06.040)
- Trabelsi D, Mhamdi R (2013) Microbial inoculants and their impact on soil microbial communities: A review. *BioMed Res Int* 11. doi:[10.1155/2013/863240](https://doi.org/10.1155/2013/863240)
- Trivedi P, Pandey A (2007) Application of immobilized cells of *Pseudomonas putida* to solubilize insoluble phosphate in broth and soil conditions. *J Plant Nutr Soil Sci* 170:629–631
- Trivedi P, Pandey A, Palni LMS (2012) Bacterial inoculants for field applications under mountain ecosystem: present initiatives and future prospects. In: Maheshwari DK (ed) *Bacteria in agrobiology: plant probiotics*. Springer-Verlag, Berlin, Heidelberg, pp. 15–44. doi:[10.1007/978-3-642-27515-9_2](https://doi.org/10.1007/978-3-642-27515-9_2)

- Veresoglou SD, Chen B, Rillig MC (2012) Arbuscular mycorrhiza and soil nitrogen cycling. *Soil Biol Biochem* 46:53–62. doi:[10.1016/j.soilbio.2011.11.018](https://doi.org/10.1016/j.soilbio.2011.11.018)
- Vinocur B, Altman A (2005) Recent advances in engineering plant tolerance to abiotic stress: achievements and limitations. *Curr Opin Biotechnol* 16:123–132. doi:[10.1016/j.copbio.2005.02.001](https://doi.org/10.1016/j.copbio.2005.02.001)
- Wagner SC (2011) Biological nitrogen fixation. *Nat Educ Knowl* 3(10):15
- Wang H, Liu S, Zhal L, Zhang J, Ren T, Fan B, Liu H (2015) Preparation and utilization of phosphate biofertilizers using agricultural waste. *J Integr Agric* 14(1):158–167. doi:[10.1016/S2095-3119\(14\)60760-7](https://doi.org/10.1016/S2095-3119(14)60760-7)
- White PJ, Karley AJ (2010) Potassium. In: Hell R, Mendel RR (eds) *Cell biology of metals and nutrients, plant cell monographs*, vol 17. Springer, Berlin, pp. 199–224
- Wu HS, Yang XM, Fan JQ, Miao WG, Ling N, Xu YU, Huang QC, Shen Q (2009) Suppression of Fusarium wilt of watermelon by a bio-organic fertilizer containing combinations of antagonistic microorganisms. *BioControl* 54:287–295. doi:[10.1007/s10526-008-9168-7](https://doi.org/10.1007/s10526-008-9168-7)
- Wu Q-S, Srivastava AK, Zou Y-N (2013) AMF-induced tolerance to drought stress in citrus: a review. *Sci Hortic* 164:77–87. doi:[10.1016/j.scienta.2013.09.010](https://doi.org/10.1016/j.scienta.2013.09.010)
- Xiao K, Kinkel LL, Samac DA (2002) Biological control of Phytophthora root rots on alfalfa and soybean with *Streptomyces*. *BioControl* 23:285–295. doi:[10.1006/Bcon.2001.1015](https://doi.org/10.1006/Bcon.2001.1015)
- Xue C, Penton CR, Shen Z, Zhang R, Huang Q, Li R, Ruan Y, Shen Q (2015) Manipulating the banana rhizosphere microbiome for biological control of Panama disease. *Sci Report* 5:11124. doi:[10.1038/srep11124](https://doi.org/10.1038/srep11124)
- Yadav J, Verma JP (2014) Effect of seed inoculation with indigenous *Rhizobium* and plant growth promoting rhizobacteria on nutrients uptake and yields of chickpea (*Cicer arietinum* L). *Eur J Soil Biol* 63:70–77. doi:[10.1016/j.ejsobi.2014.05.001](https://doi.org/10.1016/j.ejsobi.2014.05.001)
- Yang J, Kloepper J, Ryu C (2009) Rhizosphere bacteria help plants tolerate abiotic stress. *Trends Plant Sci* 14:1–4. doi:[10.1016/j.tplants.2008.10.004](https://doi.org/10.1016/j.tplants.2008.10.004)
- Yang XM, Chen LH, Yong XY, Zhana FG, Ran W, Shen QR (2011) Formulations can affect colonization and biocontrol efficiency of *Trichoderma harzianum* SQR-T037 against Fusarium wilt of cucumbers. *Biol Fertil Soils* 47:239–248. doi:[10.1007/s00374-010-0527-z](https://doi.org/10.1007/s00374-010-0527-z)
- Yousaf S, Afzal M, Reichenauer TG, Brady CL, Sessitsch A (2011) Hydrocarbon degradation, plant colonization and gene expression of alkane degradation genes by endophytic *Enterobacter ludwigii* strains. *Environ Pollut* 159(10):2675–2683. doi:[10.1016/j.envpol.2011.05.031](https://doi.org/10.1016/j.envpol.2011.05.031)
- Yuste JC, Peñuelas J, Estiarte M, Garcia-mas J, Mattana S, Ogaya R, Pujol M, Sardan J (2011) Drought-resistant fungi control soil organic matter decomposition and its response to temperature. *Glob Chang Biol* 17:1475–1486. doi:[10.1111/j.1365-2486.2010.02300.x](https://doi.org/10.1111/j.1365-2486.2010.02300.x)
- Zaidi A, Khan MS, Ahmed E (2014) Microphos principles production and application strategies. In: Khan MS, Zaidi A, Musarrat J (eds) *Phosphate solubilizing microorganism*. Springer Cham Heidelberg, New York, pp 1–30
- Zarjani JK, Aliasgharzad N, Oustan S, Emadi M, Ahmadi A (2013) Isolation and characterization of potassium solubilizing bacteria in some Iranian soils. *Arch Agron Soil Sci* 59(12):1713–1723. doi:[10.1080/03650340.2012.756977](https://doi.org/10.1080/03650340.2012.756977)
- Zhu F, Qu L, Hong X, Sun X (2011) Isolation and characterization of a phosphate-solubilizing halophilic bacterium *Kushneria* sp. YCWA18 from Daqiao Saltern on the coast of yellow sea of China. *Evid Based Complement Alternat Med*:6. doi:[10.1155/2011/615032](https://doi.org/10.1155/2011/615032)

Index

A

Abdelmageed, A.H.A., 50
 Abida, S., 97–100
 Agrawal, S.B., 115–147
 Agricultural sustainability, 282, 284
Agrobacterium rhizogenes, 214, 235, 238
 Agroforestry, 116, 310, 315, 319
 Ahmad, A., 57–108
 Ahmad, M., 99, 102
 Ahmad, T., 73
 Akbar, H., 99, 102
 Akiyama, T., 192
 Alamgir, A., 99
 Ali, K., 57–108
 Allen, D.J., 141
 Alori, E.T., 281–300
 Altman, A., 293
 Amonette, J.E., 70
 Anderson, J.A., 118
 Animal waste, 1–10, 60, 61, 73
 Aon, M., 57–108
 Arbuscular mycorrhizal fungi, 16, 17, 25–28,
 102, 117, 281, 285, 287–291, 294, 300
 Arif, M., 57–108
 Ashcroft, W.J., 257
 Azeem, M., 73, 77

B

Babalola, O.O., 281–300
 Bagge, E., 7
 Bailey, B., 41
 Baille, A., 41
 Balanced fertilization, 256, 271
 Balwinder-Singh, 256, 259, 260
 Banedjschafie, S., 271

Bano, A., 141
 Bartzanas, T., 43
 Basso, A.S., 76
 Bhagat, R.M., 263
 Biochar, 6, 57–108
 Biochar technology, 81–89, 106, 107
 Biocontrol, 194, 282–284, 295
 Biofertilizer, 147, 282, 284, 287, 294
 Biofuels, 9, 80, 213–239
 Bioremediation, 216, 219, 220, 222, 223, 283
 Biotechnology, 239
 Biswas, B.C., 270
 Borrel, A.K., 266
 Boulard, T., 43–45
 Bouman, B.A.M., 266
 Bournet, P., 45
 Bridgwater, A.V., 80
 Bunemann, E., 24

C

Caba, J.M., 142
 Cabangon, R.J., 269
 Caldwell, M.M., 137
 Chakraborty, D., 261
 Chan, K.Y., 69, 77
 Charbonneau, G.A., 142
 Chaudhary, B.U., 266, 269
 Chimphango, S.B.M., 135
 Chitinase, 163–194, 295
 Choudhary, K.K., 115–147
 Choudhary, M.A., 268, 269
 Chouhan, S., 135
 Climate change, 73, 104, 106, 118, 290, 310,
 314, 318, 320
 Conn, V.M., 299

Cooper, P.J.M., 255
 Cossu, R., 1–10
 Crop yields, 19, 29, 73–76, 99–103, 107, 108,
 255, 256, 268, 270, 271, 311, 312,
 314–319, 321

D

da Silva, A.L.L., 213–239
 Dana, M.M., 193
 Dare, M.O., 281–300
 Dempster, D.N., 77
 Direct seeded rice, 256, 266, 269–270
 Diversity, 17, 23–29, 215, 299, 300
 Dryland agriculture, 72–80
 Duba, L.I., 145
 Dullaart, J., 145

E

Environment, 3, 8, 10, 18, 29, 30, 38, 40, 45,
 47, 51, 52, 80, 102, 132, 216, 221–224,
 233, 236, 239, 260, 262, 269, 270, 282,
 286, 300, 317–319, 321
 Environmental services, 319
 Erenstein, O., 264

F

Fan pad evaporative cooling, 42, 46–52
 Fan, T., 271
 Farooq, N.I., 100, 106
 Fatnassi, H., 44, 45
 Feedstock, 58–71, 77, 81–89, 96, 103–108,
 237
 Flint, S.D., 137
 Food security, 254, 312, 314, 317, 318
 Franco, C.M.M., 299
 Fungal resistance, 185, 188

G

Gajri, P.R., 271
 Gaskin, J.W., 69–70
 Ghuman, B.S., 260
 Gill, B.S., 260
 Gill, M.S., 271
 Giroto, F., 1–10
 Glick, B.R., 294
 Glucanase, 163–194
 Gonzalez-Real, M.M., 41
 Greenhouse microclimate, 38–42, 46, 48, 49, 52
 Gruda, N., 50
 Gupta, R.K., 265

H

Habibi, P., 213–239
 Hadda, M.S., 253–272
 Harper, J.E., 141
 He, J., 139
 He, X., 309–322
 Hefferon, K., 213–239
 Herschkovitz, Y., 300
 Hira, G.S., 257
 Hobbs, P., 265
 Hughes, J.B., 220
 Hussain, Q., 57–108

I

Insect, 6, 9, 42, 45, 48, 134, 174, 218, 228
 Ippolito, J.A., 76, 77
 Izaguirre, M.M., 140

J

Jalota, S.K., 260
 Jansen, M.A.K., 140
 Jat, M.L., 264, 268
 Jha, Y., 293
 Jospeh, S., 70

K

Kameyama, K., 76
 Katyal, J.C., 270
 Kaur, B., 268
 Khan, A., 97–100
 Khan, M.J., 97–100
 Khan, N., 99
 Khan, T.S., 98
 Kittas, C., 43, 44, 46
 Kostina, E., 138
 Kubicek, C.P., 175
 Kukal, S.S., 258

L

Laing, M., 37–52
 Laird, D.A., 76
 Laser land leveling, 256, 268–269
 Latif, S., 57–108
 Laube, J.C., 118
 Legumes, 101, 116, 117, 132–146, 235, 286,
 294, 316
 Lehman, R.M., 15–30
 Lehmann, J., 97
 Leifheit, E.F., 290
 Li, S., 46, 49

Li, X., 64
 Lin, L., 8
 Liu, C.A., 271
 Liu, X.H., 65
 Lopez, A., 46
 Lou, Y., 118, 132

M

Maboko, M.M., 41, 48–50
 Mahimairaja, S., 65
 Makhzoum, A., 213–239
 Malik, R.K., 265
 Malik, S., 213–239
 Management, 3–9, 16, 17, 20, 23, 27–30, 41, 42, 58, 63, 73, 103, 116, 134, 147, 214, 234, 253–272, 286, 288, 299, 300, 310, 315, 316, 318
 Maria Fatima Grossi De Sa, 213–239
 Mashonjowa, E., 50
 Max, F.J., 48, 49
 Mian, I.A., 57–108
 Microbial inoculants, 281–300
 Mkonda, M.Y., 309–322
 Mohammad, D., 99
 Mulching, 259–262, 271, 272, 310, 314, 316–319
 Mycorrhiza, 116, 117, 145, 147, 235, 284, 288, 290

N

Naeem, M.A., 64–70
 Nair, S., 293
 Natural ventilation, 42–45, 47–51
 Nelissen, V., 64
 Newcomb, W., 142
 Newton, L., 9
 Noorollahi, Y., 8
 Novak, J.M., 77

O

Odindo, A., 37–52
 Organic fertilization, 73, 318
 Oshita, K., 8
 Oz, H., 46

P

Pakistan, 57–108
 Pan, G., 57–108
 Passioura, J., 255
 Paul, D., 293

Phosphorus, soil microorganisms, 16–30
 Phytoremediation, 213–239, 285, 291, 292
 Plant growth, 16, 22, 24–26, 28, 38, 42, 69, 80, 104, 117, 130, 131, 136, 137, 190, 191, 222, 282–284, 286–289, 291, 293, 294, 296, 300
 Plant growth promoting microorganisms (PGPM), 286
 Plant hormones, 141–146, 282, 293
 Plant molecular pharming, 228–234, 238, 239
 Planting basins, 318
 Prihar, S.S., 259
 Probanza, A., 299
 Pyrolysis, 6, 59, 60, 63–71, 76, 80–93, 95, 96, 103–105, 108

Q

Qayyum, M.F., 57–108
 Qin, J., 261
 Quilliam, 102

R

Raised bed planting, 265–268
 Rasul, F., 57–108
 Recombinant proteins, 228–233
 Relative humidity, 38, 41–48, 50, 51
 Rezek, J., 223
 Rhizobia, 116–119, 135, 144–147, 235, 284, 286, 287
 Rillig, M.C., 290
 Rodrigues, G.C., 140

S

Saghir, M., 57–108
 Sajida, P., 97–100
 Sakrabani, R., 57–108
 Sandhu, B.S., 257, 258, 266
 Sandhu, J.S., 163–194
 Sandhu, S.S., 253–272
 Santos, I., 138, 139
 Sarkar, A.A., 257
 Sasaki, C., 168
 Sauer, T.J., 260
 Secondary metabolites, 194, 214, 215, 224–228, 237, 295
 Sela-Buurlage, M.B., 168, 184
 Semi-arid, 37–52, 69, 73–75, 77, 80, 95, 317, 318
 Shackley, S., 57–108
 Shah, H., 100, 106
 Sharma, P.K., 263

- Shen, J.B., 271
 Shenbagavalli, S., 65
 Siddique, K.H.M., 260
 Sidhu, H.S., 260
 Sidhu, M.K., 163–194
 Simpson, R.J., 20
 Singh, A., 132
 Singh, B., 66
 Singh, D., 253–272
 Singh, J., 253–272
 Singh, K.B., 263
 Singh, R.P., 264
 Smith, J.L., 77
 Soccol, C.R., 213–239
 Sohi, S., 64
 Soil fertility, 16, 22, 27–30, 58, 73, 92, 103,
 105, 107, 119, 134, 270, 271, 286–289,
 310, 311, 313–317
 Soil fertility and health, 282–300
 Soni, P., 44
 Streubel, J.D., 76, 77
 Su, Z., 267
 Sudhir-Yadav, 270
 Sur, H.S., 260
 Sustainability, 107, 254, 282, 284, 289
 Symbiosis, 26, 116, 118, 132–147
 Szögi, A.A., 7
- T**
 Taheri, W.I., 15–30
 Takakura, Y., 184
 Takeuchi, A., 141
 Tanzania, 309–322
 Teitel, M., 43, 44, 47–49
 Temperature, 23–25, 38–52, 60, 64–71,
 80–89, 95, 106, 108, 236, 260, 261,
 296, 310, 314
 Tensiometer, 258
 Terraces, 310, 316, 319
- Tewatia, R.K., 270
 Thiye, E.L., 37–52
 Transgenic plants, 164, 184–190, 192,
 193, 227–229, 232, 233, 235,
 237, 239
 Transgenic roots, 235, 239
 Tuong, T.P., 266
- U**
 Ultraviolet-B (UV-B), 115–147
- V**
 Van Aalten, D.M.F., 168
 Van den Noort, R.E., 140
 Vinocur, B., 293 von
 Elsner, B., 41 von
 Zabeltitz, C., 41
- W**
 Water productivity, 254–261, 263–265,
 268–271
 Willits, D.H., 46, 49
 Workneh, T., 37–52
 Wu, H., 8
 Wulff, A., 138
- Y**
 Yadav, I.S., 163–194
 Yuste, J.C., 290
- Z**
 Zhang, J.H., 64
 Zhang, X.C., 65
 Zhu, F.X., 9
 Zhu, K., 6