

Page 1 of 4

Albanian j. agric. sci. 2017;16 (1): 10-13 Agricultural University of Tirana

*Corresponding author: Oluyemisi Fawole; E-mail: yemisifawole@yahoo.com

(Accepted for publication March 20, 2017)

ISSN: 2218-2020, © Agricultural University of Tirana

RESEARCH ARTICLE (Open Access)

Impact of Chemical Inputs on Arbuscular Mycorrhizal Spores in Soil :Response of AM Spores to Fertilizer and Herbicides

OLUYEMISI B. FAWOLE¹* AND ELIZABETH T. ALORI²

¹Department of Agronomy, University of Ilorin, P.M.B. 1515, Ilorin, Kwara state, Nigeria

²Crop and Soil Science Department, Landmark University, P.M.B. 1001, Omu Aran, Kwara state, Nigeria

Abstract

Effects of chemical inputs on abundance of arbuscular mycorrhizal (AM) fungi spores were investigated under a maize monoculture in Southern Guinea Savanna ecological zone of Nigeria. The experimental plots received 0, 60, 120 and 180 kg N/ha of fertilizer (N.P.K. 15: 15: 15) and 0, 1, 2 and 3 kg active ingredient (a.i) pre-emergence herbicide (atrazine) per hectare. AM fungal spores were isolated from the conventionally tilled, treated soils, cropped with maize by wet sieving and decanting method. Isolated AM spores were counted and spore abundance calculated. The spores were morphotyped and 13 AM fungi taxa were identified. These belong to six genera of AM fungi: *Glomus*, *Paraglomus*, *Acaulospora*, *Entrophospora*, *Gigaspora* and *Scutellospora*. Increasing atrazine concentrations resulted in significant deduction in AMF spores while increasing N.P.K. content led to an initial significant increase in AMF spores but a later decline in spore population. Both fertilization and herbicide application resulted in a decrease in AMF spore numbers at maturity of maize crop. Application of 160kg N/ha and 3kg a.i. ha⁻¹ of atrazine caused a significant reduction in spore density (117g⁻¹ of soil as against 202 g⁻¹ of soil at 0 kg N ha⁻¹ and 0kg a.i. ha⁻¹ atrazine).

Keywords: Arbuscular Mycorrhizal Spores, Herbicide, Fertilizer, Southern Guinea Savanna

1. Introduction

Arbuscular Mycorrhizal Fungi (AMF) are obligate biotrophs that form symbiotic interactions with a wide range of plants, wild and cultivated plants alike. They are ubiquitous in natural and agricultural soil [8]. AMF improve plant mineral nutrition thereby playing important role in nutrient cycling plant health and plant water relations [1, 5]. Atrazine is one of the most widely used agricultural herbicide worldwide. It may be applied before and after planting to control broad leaf and grassy weeds. Atrazine is applied on maize, sorghum and sugarcane.

A major problem facing crop production in Southern Guinea Savanna ecological zone of Nigeria is the low fertility status of the soil [4]. This therefore necessitates the use of large dressing of inorganic fertilizer for maximum plant growth and yield. We hypothesized that agricultural inputs (Inorganic fertilizer and herbicide) would affect the population density of the AM fungal community. Loss of AMF species could lead to irreversible destruction of habitat and eventual loss of agroecosystem. There is little information on the effect of intensive agricultural systems on AMF in Southern Guinea Savanna ecological zone of Nigeria. This research is therefore aimed at evaluating the effects of fertilizer and herbicide application on the population of AMF spores in soils cropped with maize.

2. Materials and Methods

The field experiment was carried out at the Teaching and Research Farm of the University of Ilorin, Ilorin, Nigeria. The soil examined was an Alfisol with the following composition (at 0-15 cm layer) % Organic Matter 0.48, %N 210, P (mg kg⁻¹) 1.88 and soil pH was 5.85. The agricultural inputs were herbicide (Atrazine 50% Sc.) treatment: 0, 1, 2 and 3kg active ingredient per hectare applied as pre-emergence; fertilization treatment (N. P. K. 15:15:15) : 0 Kg ha⁻¹ N, 60 Kg ha⁻¹ N, 12 Kg ha⁻¹ N 0 and 180 Kg ha⁻¹ N at two weeks after planting. The test crop was maize. The treatments were arranged in randomized split plot design and replicated three times. To study AM fungal

Page 2 of 4

Fawole O. B. and E. T. Alori, 2017

spore population soil samples were collected at 4weeks after planting and at maturity. AMF community were studied by spore extraction from soil. Hence 100g soil (dry weight) of each soil sample was wet sieved and centrifuged, following the methodology proposed by [2]. Quantification was carried out in 9cm diameter petri dish with a grid line of 1cm per side

under stereoscopic microscope at X40. Ten divisions were counted and related to total number of spores by using the method modified by Mckenney and Lindsey [6]. Data collected were subjected to analysis of variance. Means were separated using Least Significant Difference (LSD).

3. Results and Discussion

Thirteen (13) AMF taxa belonging to 6 genera of AMF: *Glomus*, *Paraglomus*, *Acaulospora*, *Entrophospora*, *Gigaspora* and *Scutellospora* were identified from the study site. The percentage frequencies of the species recorded are shown in (fig.1). The effects of fertilizer (N. P. K 15: 15: 15) and herbicide (Atrazine) on the population of AMF spores are shown in Tables 1 and 2. A significant reduction in AMF spores associated with the increase in atrazine concentration agrees with the findings that fumigation to destroy weeds also kills AMF [7]. Increasing N. P. K. content led to significant increase in AMF spores at 4 weeks after planting (WAP) (Table 1), but an increase in the quantity of both herbicide and fertilizer used resulted in a significant decrease in the population of AMF spores at maturity of test plant. Application of 160kg N/ha and 3kg a.i/ha of atrazine caused a significant reduction in spore density (117/g of soil as against 202/g of soil at 0kg N/ha and 0kg ai/ha atrazine). This suggests that increase in fertilizer application probably supported an increase in AMF spore numbers until a critical level where a further increase in the availability of inorganic fertilizer in soil resulted in a decrease in number of AMF spores. This is in line with the findings reported in [8] who stated that fertilizer and pesticide applications to soil may counteract super optimum AM fungal populations. Reduction in mycorrhizal population will have adverse effects on mycorrhizal infections and uptake of nutrients by mycorrhizal plants. It was also reported that application of atrazine resulted in a decrease in AMF colonization [3]. Elimination of AMF populations can lead to problems with plant establishment and survival. Nitrogen additions decrease investment in AMF by plants (assessed primarily as % Root Length Colonization (RLC)) by an average of 24% in field studies [10]. Figure 1. % Frequency of occurrence of AMF isolates. Bars are % error

0
2
4
6
8
10
12
14
16

18

Frequency of Occurrence (%)

AMF Isolates

11

Page 3 of 4

Impact of chemical inputs on arbuscular mycorrhizal spores in soil

Table 1. Interactive effect of herbicide (Atrazine) and fertilizer (N.P.K) on AMF population at 4WAP

HERBICIDE

FERTILIZER H1 H2 H3 H4 MEAN

F1 534e 505g 485h 421k 486

F2 545d 540de 508g 443j 509

F3 576b 556c 525f 478i 534

F4 613a 579b 542d 506g 560

MEAN 567 545 515 462 522

LSD 2.98

1Means followed by the same letter (s) in column and row of any set of treatment are not significantly different at

5% level of probability using least significant difference (LSD) test

Table 2: Effect of herbicide (Atrazine) and fertilizer (N.P.K) on the AMF spore population

TREATMENT AMF SPORE POPULATION

HERBICIDE (H) 4WAP MATURITY

H1 567a 437a

H2 545b 337b

H3 515c 228c

H4 462d 155d

MEAN 522.25 289.25

LSD 1.81 4.32

FERTILIZER (F)

F1 486d 355a

F2 509c 304b

F3 534b 267c

F4 560a 231d

MEAN 522 289

LSD 1.49 3.35

INTERACTIVE (H & F) 2.98 6.70

1Means followed by the same letter (s) in column of any set of treatment are not significantly different at 5% level of probability

using least significant difference (LSD) test

2Key: F1=Fertilizer at 0kg, N/ha, F2=Fertilizer at 80Kg N/ha, F3=120Kg N/ha, F4=160Kg N/ha, H1=herbicide at 0 kg ai/ha,

H2=herbicide at 1 kg a.i/ha, H3=herbicide at 2 kg a.i/ha, H4=herbicide at 3 kg a.i/ha.

4. Conclusions

The results of this study suggest that

conventional agriculture that relies solely on inputs

such as mineral fertilizers & herbicides for increasing

productivity do reduce the population of AM spores in

soil and would thus have a long term negative impact

on mycorrhizal infections. There is therefore the need

for use of more eco-friendly soil inputs for sustainable

crop production.

5. References

1. Bonfante, P. & Genre, A: Arbuscular mycorrhizal dialogues: do you speak 'plantish' or 'fungish'? Trends in Plant Science 2014, 150 [3]: 150-4.
2. Brundrett, M., Bougher, N., Dell, B., Grove, T. & Malajczuk, N: Working with mycorrhizas in forestry and agriculture, Australian Centre for international Agricultural Research 1996.
3. Demir, S., Celebi, S. Z., Kaya, I. & Sahar, A. K: The applications of arbuscular mycorrhizal fungus (AMF) and atrazine on silage maize 12

Page 4 of 4

Fawole O. B. and E. T. Alori, 2017 (Zea mays L.) at different irrigation regimes.

Journal of Food Agriculture and Environment 2011, 9:629-633.

4. Fawole, O. B. & Olowonihi, E. T: Studies on the Distribution of Bacteria and Fungi in the five soil series of University of Ilorin Teaching and Research Farm. Journal of Agricultural Research and Development 2005, 4: 218-227.

5. Li, T., Lin, G., Zhang, X., Chen, Y. L., Zhang, S.

B. & Chen, B. D: Relative importance of an arbuscular mycorrhizal fungus (Rhizophagus intraradices) and root hairs in plant drought tolerance. Mycorrhiza 2014, 24: 595-602.

6. Mckenney, M. C. & Lindsey, D. L: Improved method of quantifying endomycorrhizal spores from soil. *Mycologia* 1987, 79:779-782.
7. Owusu-Bennoah E:. Mycorrhizal in tropical agriculture. *Ghana Journal of agricultural science* 1995, 24-27, 11-18.
8. Sharif M. & Moawad A.M: Arbuscular Mycorrhiza Incidence and Infectivity of Crops in Northwest Frontier Province of Pakistan. *World J. Agric Sci.* 2006, 2: 123-132.
9. Smith, F. A. & Smith, S. E: What is the significance of the arbuscular mycorrhizal colonisation of many economically important crop plants? . *Plant and Soil* 348: 63-79.
10. Treseder, K. K: A meta-analysis of mycorrhizal responses to nitrogen, phosphorus, and atmospheric CO₂ in field studies *New Phytologist* 2004, 164: 347-355.

13



3_FAWOLE 9-13 pp.pdf

3_FAWOLE 9-13 pp.pdf

Sign In

Page 1 of 4